

Ref #	Hits	Search Query	DBs	Default Operator	Plurals	Time Stamp
L1	289	hybrid\$ near20 nanoparticle	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2006/01/27 17:44
L2	34071	435/6[ccls]	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2006/01/27 17:44
L3	164	I1 and I2	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2006/01/27 17:44
L4	1107147	@rlad<"20030227"	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2006/01/27 17:44
L5	112	I3 and I4	US-PGPUB; USPAT; EPO; JPO; DERWENT; IBM_TDB	OR	ON	2006/01/27 17:44

\*\*\*\*\* Welcome to STN International \*\*\*\*\*  
\*\*\*\*\*

NEWS 1 Web Page URLs for STN Seminar Schedule - N. America  
NEWS 2 "Ask CAS" for self-help around the clock  
NEWS 3 DEC 05 CASREACT(R) - Over 10 million reactions available  
NEWS 4 DEC 14 2006 MeSH terms loaded in MEDLINE/LMEDLINE  
NEWS 5 DEC 14 2006 MeSH terms loaded for MEDLINE file segment of TOXCENTER  
NEWS 6 DEC 14 CA/CAPLUS to be enhanced with updated IPC codes  
NEWS 7 DEC 21 IPC search and display fields enhanced in CA/CAPLUS with the IPC reform  
NEWS 8 DEC 23 New IPC8 SEARCH, DISPLAY, and SELECT fields in USPATFULL/USPAT2  
NEWS 9 JAN 13 IPC 8 searching in IFIPAT, IFIUDB, and IFICDB  
NEWS 10 JAN 13 New IPC 8 SEARCH, DISPLAY, and SELECT enhancements added to INPADOC  
NEWS 11 JAN 17 Pre-1988 INPI data added to MARPAT  
NEWS 12 JAN 17 IPC 8 in the WPI family of databases including WPIFV

NEWS EXPRESS JANUARY 03 CURRENT VERSION FOR WINDOWS IS V8.01, CURRENT MACINTOSH VERSION IS V6.0c(ENG) AND V6.0j(JP), AND CURRENT DISCOVER FILE IS DATED 19 DECEMBER 2005. V8.0 USERS CAN OBTAIN THE UPGRADE TO V8.01 AT  
<http://download.cas.org/express/v8.0-Discover/>

NEWS HOURS STN Operating Hours Plus Help Desk Availability  
NEWS INTER General Internet Information  
NEWS LOGIN Welcome Banner and News Items  
NEWS PHONE Direct Dial and Telecommunication Network  
Access to STN  
NEWS WWW CAS World Wide Web Site (general information)

Enter NEWS followed by the item number or name to see news on that specific topic.

All use of STN is subject to the provisions of the STN Customer agreement. Please note that this agreement limits use to scientific research. Use for software development or design or implementation of commercial gateways or other similar uses is prohibited and may result in loss of user privileges and other penalties.

\*\*\*\*\* STN Columbus \*\*\*\*\*  
\*\*\*\*\*

FILE 'HOME' ENTERED AT 17:50:38 ON 27 JAN 2006

=> file caplus	
COST IN U.S. DOLLARS	SINCE FILE
TOTAL	ENTRY SESSION
FULL ESTIMATED COST	0.21 0.21

FILE 'CAPLUS' ENTERED AT 17:50:48 ON 27 JAN 2006  
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.  
COPYRIGHT (C) 2006 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 27 Jan 2006 VOL 144 ISS 6  
FILE LAST UPDATED: 26 Jan 2006 (20060126/ED)

Effective October 17, 2005, revised CAS Information Use Policies apply.  
They are available for your review at:

<http://www.cas.org/infopolicy.html>

=> s (hybridi? (20a) nanoparticle#)/bi,ab 168171 HYBRIDI?/BI  
144210 HYBRIDI?/AB 56078 NANOPARTICLE#/BI  
38709 NANOPARTICLE#/AB  
L1 373 (HYBRIDI? (20A) NANOPARTICLE#)/BI,AB

=> l1 and (rna# or dna# or cdna# or mrna#)/bi,ab  
L1 IS NOT A RECOGNIZED COMMAND  
The previous command name entered was not recognized by the system.  
For a list of commands available to you in the current file, enter "HELP COMMANDS" at an arrow prompt (=>).

=> s l1 and (rna# or dna# or cdna# or mrna#)/bi,ab 306951  
RNA#/BI 234351 RNA#/AB 758681  
DNA#/BI 591971 DNA#/AB 194569  
CDNA#/BI 152927 CDNA#/AB 284305  
MRNA#/BI 249121 MRNA#/AB  
L2 305 L1 AND (RNA# OR DNA# OR CDNA# OR MRNA#)/BI,AB

=> s l2 not 2006/py 73774 2006/PY  
L3 291 L2 NOT 2006/PY

=> s l3 not 2005/py 1188235 2005/PY  
L4 193 L3 NOT 2005/PY

=> s l4 not 2004/py 1284082 2004/PY  
L5 120 L4 NOT 2004/PY

=> s l5 not 2003/py 1257963 2003/PY  
L6 63 L5 NOT 2003/PY

=> d his  
(FILE 'HOME' ENTERED AT 17:50:38 ON 27 JAN 2006)  
FILE 'CAPLUS' ENTERED AT 17:50:48 ON 27 JAN 2006  
L1 373 S (HYBRIDI? (20A) NANOPARTICLE#)/BI,AB  
L2 305 S L1 AND (RNA# OR DNA# OR CDNA# OR MRNA#)/BI,AB  
L3 291 S L2 NOT 2006/PY

L4 193 S L3 NOT 2005/PY  
L5 120 S L4 NOT 2004/PY  
L6 63 S L5 NOT 2003/PY

=> d l6 1-63 bib ab

L6 ANSWER 1 OF 63 CAPLUS COPYRIGHT 2006 ACS on STN  
AN 2003:168705 CAPLUS  
DN 138:349216  
TI Sensing and diagnosis  
AU Maeda, Mizuo  
CS Graduate School of Engineering, Kyushu University,  
Fukuoka, Japan  
SO Supramolecular Design for Biological Applications (2002),  
287-298. Editor(s): Yui, Nobuhiko. Publisher: CRC Press LLC,  
Boca Raton, Fla. CODEN: 69DQPD; ISBN: 0-8493-0965-4  
DT Conference  
LA English

AB Two types of supramol. assembling phenomena are discussed in relation to gene mutation assays: \*\*\*DNA\*\*\* - carrying colloidal nanoparticles prep. from \*\*\*DNA\*\*\* - PNIPAAm graft copolymer through supramol. assembly; and \*\*\*DNA\*\*\* -carrying nanoparticles assembled in the presence of complementary \*\*\*DNA\*\*\* to yield visibly detectable aggregates due to crosslinking or colloidal stability change. A typical example of supramol. systems, useful for biol. sensing and diagnosis, is explained. The author proposes a novel aggregation mechanism of \*\*\*DNA\*\*\* -contg. colloidal particles, which is based on the stability decrease of colloidal particles accompanied by the duplex formation of the shell \*\*\*DNA\*\*\* with the complementary \*\*\*DNA\*\*\*. Using the particle aggregation assay, sequence- and chain length-selective \*\*\*DNA\*\*\* detection is attained.

RE.CNT 28 THERE ARE 28 CITED REFERENCES AVAILABLE FOR THIS RECORD ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 2 OF 63 CAPLUS COPYRIGHT 2006 ACS on STN  
AN 2003:149917 CAPLUS  
DN 138:215997

TI Bioconjugated nanoparticle probes for ultrasensitive \*\*\*DNA\*\*\* detection

AU Taylor, Jason Roger  
CS Indiana Univ., Bloomington, IN, USA  
SO (2002) 187 pp. Avail.: UMI, Order No. DA3054502 From: Diss. Abstr. Int., B 2002, 63(5), 2363  
DT Dissertation  
LA English  
AB Unavailable

L6 ANSWER 3 OF 63 CAPLUS COPYRIGHT 2006 ACS on STN  
AN 2003:24393 CAPLUS  
DN 138:50796

TI Nanoparticle-labeled probe and uses in \*\*\*DNA\*\*\* chip for diagnosis  
IN Pang, Daiwen; Wang, Yefu; Zhang, Zhiling; Cao, Junping; Cai, Ruxiu; Zheng, Guzhi  
PA Wuhan Univ., Peop. Rep. China  
SO Faming Zhuanli Shenqing Gongkai Shuomingshu, 16 pp. CODEN: CNXXEV

DT Patent

LA Chinese

FAN.CNT 1 PATENT NO. KIND DATE APPLICATION  
NO. DATE -----

PI CN 1339609 A 20020313 CN 2001-133527  
20010930  
PRAI CN 2001-133527 20010930

AB The invention relates to prepn. of nanoparticle-labeled probe of \*\*\*DNA\*\*\* chip for diagnosis. The said nanoparticle-labeled probe is the 100-1,000 bp single-stranded \*\*\*RNA\*\*\* or \*\*\*DNA\*\*\*, which were labeled with Au, Ag, CdS, Fe2O3, or SiO2. The \*\*\*nanoparticle\*\*\* -labeled probe is immobilized on glass, nylon membrane or cellulose sulfate membrane and \*\*\*hybridized\*\*\* in \*\*\*DNA\*\*\* chip. The hybridization signals are detected by silver staining using Ag salt and reductant.

L6 ANSWER 4 OF 63 CAPLUS COPYRIGHT 2006 ACS on STN  
AN 2002:861903 CAPLUS  
DN 138:251030

TI Fabrication of two- and three-dimensional structures of \*\*\*nanoparticles\*\*\* using LB method and \*\*\*DNA\*\*\* \*\*\*hybridization\*\*\*

AU Takahagi, Takayuki; Huang, Shujuan; Tsutsui, Gen; Sakaue, Hiroyuki; Shingubara, Shoso

CS Graduate School of Advanced Sciences of Matter, Hiroshima University, Higashi-Hiroshima, 739-8526, Japan

SO Materials Research Society Symposium Proceedings (2002), 707(Self-Assembly Processes in Materials), 87-92 CODEN: MRSPDH; ISSN: 0272-9172

PB Materials Research Society

DT Journal

LA English

AB In this paper we describe fabrication methods for two types of nanostructures, two- and three-dimensional arrays of gold nanoparticles. Large-scale and high-ordered monolayers of alkanethiol-encapsulated gold particles were fabricated by using Langmuir-Blodgett (LB) method. Three-dimensional nanoparticle arrays composed of gold \*\*\*nanoparticles\*\*\* of two different sizes, which were encapsulated by complementary thiol-capped \*\*\*DNA\*\*\* oligonucleotides, were fabricated by using \*\*\*DNA\*\*\* \*\*\*hybridization\*\*\*. \*\*\*DNA\*\*\* hybridization occurred upon mixing these particles, which resulted in the assembly of three-dimensional nanostructure of gold particles. SEM observations and UV spectroscopy measurement were performed to confirm the construction of the nanostructures.

RE.CNT 9 THERE ARE 9 CITED REFERENCES AVAILABLE FOR THIS RECORD ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 5 OF 63 CAPLUS COPYRIGHT 2006 ACS on STN  
AN 2002:818318 CAPLUS  
DN 138:33867

TI Electrochemical detection of \*\*\*DNA\*\*\* \*\*\*hybridization\*\*\* based on silver-enhanced gold \*\*\*nanoparticle\*\*\* label

AU Cai, Hong; Wang, Yanqing; He, Pingang; Fang, Yuzhi  
CS Department of Chemistry, East China Normal University, Shanghai, 200062, Peop. Rep. China

SO Analytica Chimica Acta (2002), 469(2), 165-172 CODEN: ACACAM; ISSN: 0003-2670

PB Elsevier Science B.V.

DT Journal

LA English

AB An electrochem. detection method for analyzing sequence-specific \*\*\*DNA\*\*\* using gold nanoparticle \*\*\*DNA\*\*\* probes and subsequent signal amplification step by silver enhancement is described. The assay relies on the electrostatic adsorption of target oligonucleotides onto the sensing surface of

the glassy carbon electrode (GCE) and its \*\*\*hybridization\*\*\* to the gold \*\*\*nanoparticle\*\*\* -labeled oligonucleotides \*\*\*DNA\*\*\* probe. After silver deposition onto gold nanoparticles, binding events between probe and target were monitored by the differential pulse voltammetry (DPV) signal of the large no. of silver atoms anchored on the hybrids at the electrode surface. The signal intensity difference permits to distinguish between the match of two perfectly matched \*\*\*DNA\*\*\* strands and the near-perfect match where just one base pair was wrong. Coupled with this "nanoparticle-promoted" redn. of silver signal amplification method, the sensitivity of this electrochem. \*\*\*DNA\*\*\* biosensor has been increased by approx. two orders of magnitude and a detection limit of 50 pM of complementary oligonucleotides was obtained.  
RE.CNT 37 THERE ARE 37 CITED REFERENCES AVAILABLE FOR THIS RECORD ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 6 OF 63 CAPLUS COPYRIGHT 2006 ACS on STN  
AN 2002:818291 CAPLUS  
DN 138:20108  
TI Immobilization of oligonucleotides onto silica  
\*\*\*nanoparticles\*\*\* for \*\*\*DNA\*\*\* \*\*\*hybridization\*\*\* studies  
AU Hilliard, Lisa R.; Zhao, Xiaojun; Tan, Weihong  
CS Center for Research at Bio/Nano Interface, Department of Chemistry, and the McKnight Brain Institute, University of Florida, Gainesville, FL, 32611, USA  
SO Analytica Chimica Acta (2002), 470(1), 51-56 CODEN: ACACAM; ISSN: 0003-2670  
PB Elsevier Science B.V.  
DT Journal  
LA English  
AB This paper describes the development of oligonucleotide-functionalized nanoparticles. We used disulfide-coupling chem. for the immobilization of oligonucleotides onto silica nanoparticles and subsequently demonstrated the properties of the resulting \*\*\*DNA\*\*\* nanoparticles. Factors influencing the immobilization and hybridization processes were examd. and optimized. The oligonucleotide-modified silica \*\*\*nanoparticles\*\*\* provide an efficient substrate for \*\*\*hybridization\*\*\* and can be used in the development of \*\*\*DNA\*\*\* biosensors and biochips.  
RE.CNT 20 THERE ARE 20 CITED REFERENCES AVAILABLE FOR THIS RECORD ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 7 OF 63 CAPLUS COPYRIGHT 2006 ACS on STN  
AN 2002:791489 CAPLUS  
DN 138:34812  
TI Electrophoretic and Structural Studies of \*\*\*DNA\*\*\* - Directed Au Nanoparticle Groupings  
AU Zanchet, Daniela; Micheel, Christine M.; Parak, Wolfgang J.; Gerion, Daniele; Williams, Shara C.; Alivisatos, A. Paul  
CS Department of Chemistry, University of California, Berkeley, CA, 94720, USA  
SO Journal of Physical Chemistry B (2002), 106(45), 11758-11763 CODEN: JPCBFK; ISSN: 1520-6106  
PB American Chemical Society  
DT Journal  
LA English  
AB Discrete Au nanoparticle/ \*\*\*DNA\*\*\* conjugates have been isolated by electrophoresis and used to form small groupings of particles, such as dimers and trimers. The use of purified conjugates leads to a higher yield of the target structure, and it has allowed us better control and understanding of the

system. Newly accessible questions, such as the electrophoretic mobility of nanoparticle/ \*\*\*DNA\*\*\* hybrids and the crit. role of particle surface charge on mobility, have been studied. Detailed characterization by transmission electron microscopy (TEM) has now been done because of the higher quality of the samples. A computer program to generate pair distribution functions from TEM images was developed, pointing out the dependence of interparticle distance with \*\*\*DNA\*\*\* length on dimers of particles.  
RE.CNT 30 THERE ARE 30 CITED REFERENCES AVAILABLE FOR THIS RECORD ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 8 OF 63 CAPLUS COPYRIGHT 2006 ACS on STN  
AN 2002:691797 CAPLUS  
DN 138:83874  
TI Novel gene diagnosis using \*\*\*DNA\*\*\* conjugates  
AU Takarada, Tohru; Maeda, Mizuo  
CS Japan  
SO Kino Zairyo (2002), 22(8), 13-21 CODEN: KIZAEP; ISSN: 0286-4835  
PB Shi Emu Shi Shuppan  
DT Journal; General Review  
LA Japanese  
AB A review. The detection of gene mutation such as SNP (single nucleotide polymorphism) based on \*\*\*DNA\*\*\* \*\*\*hybridization\*\*\* using colloidal \*\*\*nanoparticles\*\*\* bearing specific \*\*\*DNA\*\*\* probes. The prepn. of the graft copolymer ( \*\*\*DNA\*\*\* -poly-N- isopropylacrylamide conjugate) and the use of the particles in the \*\*\*DNA\*\*\* hybridization anal. by turbidimetry based on aggregation of the particles were discussed. The physico-chem. principles underlying the aggregation of the particles triggered by dsDNA formation and the mechanisms of detection of the SNP by the degree of particle aggregation were discussed. The SNP anal. using the microelectrophoresis-equipped chip system and the fabrication of the \*\*\*DNA\*\*\* -conjugated affinity capillary matrix were also discussed.

L6 ANSWER 9 OF 63 CAPLUS COPYRIGHT 2006 ACS on STN  
AN 2002:689752 CAPLUS  
DN 138:33848  
TI Colorimetric SNP analysis using oligonucleotide-modified nanoparticles  
AU Ihara, Toshihiro; Chikaura, Yasushi; Tanaka, Shojiro; Jyo, Akinori  
CS Department of Applied Chemistry and Biochemistry, Faculty of Engineering, Kumamoto University, Kumamoto, 860-8555, Japan  
SO Chemical Communications (Cambridge, United Kingdom) (2002), (18), 2152-2153 CODEN: CHCOFS; ISSN: 1359-7345  
PB Royal Society of Chemistry  
DT Journal  
LA English  
AB A point mutation in the p53 gene has been detected by means of fluorescence microscopy and fluorescent resonance energy transfer (FRET) through sequence selective aggregation of \*\*\*DNA\*\*\* -modified nanoparticles, in which fluorescent dyes were impregnated.  
RE.CNT 14 THERE ARE 14 CITED REFERENCES AVAILABLE FOR THIS RECORD ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 10 OF 63 CAPLUS COPYRIGHT 2006 ACS on STN  
AN 2002:684905 CAPLUS  
DN 138:35578

TI Electrochemical stripping detection of \*\*\*DNA\*\*\*  
\*\*\*hybridization\*\*\* based on cadmium sulfide  
\*\*\*nanoparticle\*\*\* tags  
AU Wang, Joseph; Liu, Guodong; Polsky, Ronen; Merkoci, Arben  
CS Department of Chemistry and Biochemistry, New Mexico  
State University, Las Cruces, NM, 88003, USA  
SO Electrochemistry Communications (2002), 4(9), 722-726  
CODEN: ECCMF9; ISSN: 1388-2481  
PB Elsevier Science B.V.  
DT Journal  
LA English  
AB We report on the detection of \*\*\*DNA\*\*\*  
\*\*\*hybridization\*\*\* in connection to cadmium sulfide  
\*\*\*nanoparticle\*\*\* tracers and electrochem. stripping  
measurements of the cadmium. A \*\*\*nanoparticle\*\*\* -  
promoted cadmium pptn. is used to enlarge the  
\*\*\*nanoparticle\*\*\* tag and amplify the stripping \*\*\*DNA\*\*\*  
\*\*\*hybridization\*\*\* signal. In addn. to measurements of the  
dissolved cadmium ion we demonstrate solid-state measurements  
following a magnetic collection of the magnetic-bead/  
\*\*\*DNA\*\*\* -hybrid/CdS-tracer assembly onto a thick-film  
electrode transducer. The new protocol combines the  
amplification features of nanoparticle/polynucleotides assemblies  
and highly sensitive stripping potentiometric detection of  
cadmium, with an effective magnetic isolation of the duplex. The  
low detection limit (100 fmol) is coupled to good reproducibility  
(RSD=6%). Prospects for using binary inorg. colloids for multi-  
target detection are discussed.  
RE.CNT 17 THERE ARE 17 CITED REFERENCES AVAILABLE  
FOR THIS RECORD ALL CITATIONS AVAILABLE IN THE RE  
FORMAT

L6 ANSWER 11 OF 63 CAPLUS COPYRIGHT 2006 ACS on STN  
AN 2002:649353 CAPLUS  
DN 137:347094  
TI Development, characterization, and optimization of a  
\*\*\*nanoparticle\*\*\* -based \*\*\*dna\*\*\* \*\*\*hybridization\*\*\*  
detection strategy  
AU Sauthier, Marc Louis  
CS North Carolina State Univ., Raleigh, NC, USA  
SO (2001) 153 pp. Avail.: UMI, Order No. DA3027846 From:  
Diss. Abstr. Int., B 2002, 62(10), 4519  
DT Dissertation  
LA English  
AB Unavailable

L6 ANSWER 12 OF 63 CAPLUS COPYRIGHT 2006 ACS on STN  
AN 2002:638197 CAPLUS  
DN 137:180749  
TI Detection of genetic polymorphisms using generic molecular  
beacon probes labeled with fluoresce dye-conjugated metallic or  
semiconductor nanoparticles  
IN Phillips, Vince; Watson, Andrew R.; Wong, Edith  
PA USA  
SO U.S. Pat. Appl. Publ., 27 pp. CODEN: USXXCO  
DT Patent  
LA English  
FAN.CNT 1 PATENT NO. KIND DATE APPLICATION  
NO. DATE -----  
PI US 2002115082 A1 20020822 US 2001-945379  
20010831  
PRAI US 2000-230186P P 20000901  
AB Methods, compns. and articles of manuf. for assaying a  
sample for an amplification product from a target polynucleotide  
are provided. An amplification reaction is used to produce the

amplification product from the target polynucleotide so that it can  
be used to indirectly assay the sample for the target  
polynucleotide. A sample suspected of contg. the target  
polynucleotide is contacted with first and second primers to  
amplify the target polynucleotide; the first primer comprises a tag  
sequence, the complement of which is formed on the opposite  
strand during amplification and is referred to as a capture  
sequence. That opposite strand is referred to as a second primer  
extension product or an amplification product. A generic probe  
polynucleotide is provided that is a mol. beacon and can bind to  
the capture sequence to form an amplification product detection  
complex. The mol. beacon probes can be labeled with fluoresce  
dye or metallic or semiconductor nanoparticles to increase the  
sensitivity and specificity in the detection and enable  
multiplexing. Methods of detecting the amplification product  
detection complex thus produced are also provided, as are  
amplification product assay arrays, along with methods of  
forming the same. The methods are particularly useful in  
multiplex settings where a plurality of target polynucleotides are  
to be assayed. Kits comprising reagents for performing such  
methods are also provided.

L6 ANSWER 13 OF 63 CAPLUS COPYRIGHT 2006 ACS on STN  
AN 2002:614995 CAPLUS  
TI Photopatterning of \*\*\*DNA\*\*\* for integration with  
microfabricated structures  
AU Pathak, Srikant; Dentinger, Paul M.  
CS Materials Chemistry, Sandia National Laboratories,  
Livermore, CA, 94551-0969, USA  
SO Abstracts of Papers, 224th ACS National Meeting, Boston,  
MA, United States, August 18-22, 2002 (2002), COLL-006  
Publisher: American Chemical Society, Washington, D. C. CODEN:  
69CZPZ  
DT Conference; Meeting Abstract  
LA English  
AB Photopatterning of \*\*\*DNA\*\*\* on surfaces has been  
successfully used for assay development and a variety of  
research purposes. In this paper we show a facile method of  
patterning \*\*\*DNA\*\*\* with a com. proximity aligner and show  
micron-scale resolu. This method addnl. allows for alignment of  
patterned \*\*\*DNA\*\*\* to previous microfabrication steps.  
Silicon wafers were modified with monolayers of various silanes  
coupling agents. A photoacid generating polymer was then  
applied to the wafers and exposed through a Cr/Au mask.  
Several chem. strategies were then used to pattern the modified  
Si wafers using acid catalyzed coupling chem. The polymer is  
removed from the wafer and oligonucleotides are subsequently  
coupled to the surface. The most successful oligo patterning  
resulted from reacting a perfluorinated glycidyl ether to exposed  
areas of an epoxypropyl trimethoxysilane treated surface. The  
exposed areas become highly hydrophobic with reaction of the  
perfluorinated mol. to the surface. The entire wafer is then  
exposed to 5' aminated, 20 base long oligonucleotides for  
oligonucleotide attachment to the unexposed regions. The  
resulting \*\*\*DNA\*\*\* patterns were then studied for their  
binding efficiency vis-a-vis \*\*\*hybridization\*\*\* with a  
complementary strand labeled with (a) fluorescent dye; and (b)  
20 nm gold \*\*\*nanoparticles\*\*\*, and c) 20 nm gold particles  
followed by silver amplification. Stability as well as specificity of  
patterned \*\*\*DNA\*\*\* prepd. this way has been assessed in a  
no. of solvent, temp. and chem. coupling conditions. Micron-scale  
patterns along with very little -non-specific- interaction of  
complementary \*\*\*DNA\*\*\* as obsd. in our method allows one  
to use such hybrid approach towards assembly of useful  
oligonucleotide micro and nano- devices.

L6 ANSWER 14 OF 63 CAPLUS COPYRIGHT 2006 ACS on STN  
AN 2002:604378 CAPLUS  
DN 138:67154

TI SNPs analysis using \*\*\*DNA\*\*\* -modified nanoparticles  
AU Ihara, Toshihiro  
CS Department of Applied Chemistry and Biochemistry, Faculty  
of Engineering, Kumamoto University, Kumamoto, 860-8555,  
Japan

SO Bio Industry (2002), 19(8), 27-35 CODEN: BIINEG; ISSN:  
0910-6545

PB Shi Emu Shi Shuppan  
DT Journal; General Review  
LA Japanese

AB A review. The use of various nanoparticles with immobilized  
\*\*\*DNA\*\*\* mols. for SNP (single nucleotide polymorphism)  
anal. was discussed. The phase-change in surface plasmon  
polariton from the \*\*\*DNA\*\*\* -immobilized gold particles  
upon aggregation was utilized to monitor the \*\*\*DNA\*\*\*  
\*\*\*hybridization\*\*\* took place on the \*\*\*nanoparticles\*\*\*.  
Detection methods of \*\*\*DNA\*\*\* -hybridization by fluorometry  
or FRET-anal. were also discussed regarding their tech. principles  
based on avidin-biotin interaction and energy-transfer took place  
on the nanoparticles such as nanosphere. The actual example of  
exptl. results from a FRET SNP anal. for the p53 gene typing was  
presented.

L6 ANSWER 15 OF 63 CAPLUS COPYRIGHT 2006 ACS on STN  
AN 2002:602860 CAPLUS  
DN 137:266492

TI Fabrication of two- and three-dimensional structures of  
\*\*\*nanoparticles\*\*\* using LB method and \*\*\*DNA\*\*\*  
\*\*\*hybridization\*\*\*

AU Takahagi, Takayuki; Huang, Shujuan; Tsutsui, Gen; Sakaue,  
Hiroyuki; Shingubara, Shoso

CS Graduate School of Advanced Sciences of Matter, Hiroshima  
University, Higashi-Hiroshima, 739-8526, Japan

SO Materials Research Society Symposium Proceedings (2002),  
704(Nanoparticulate Materials), 47-52 CODEN: MRSPDH; ISSN:  
0272-9172

PB Materials Research Society  
DT Journal  
LA English

AB We describe fabrication methods for 2 types of  
nanostructures, 2- and 3D arrays of Au nanoparticles. Large-  
scale and high-ordered monolayers of alkanethiol-encapsulated  
Au particles were fabricated by Langmuir-Blodgett (LB) method.  
Three-dimensional nanoparticle arrays composed of Au  
\*\*\*nanoparticles\*\*\* of 2 different sizes, which were  
encapsulated by complementary thiol-capped \*\*\*DNA\*\*\*  
oligonucleotides, were fabricated by \*\*\*DNA\*\*\*  
\*\*\*hybridization\*\*\*. \*\*\*DNA\*\*\* hybridization occurred  
upon mixing these particles, which resulted in the assembly of 3D  
nanostructure of Au particles. SEM observations and UV  
spectroscopy measurement were performed to confirm the  
construction of the nanostructures.

RE.CNT 9 THERE ARE 9 CITED REFERENCES AVAILABLE FOR  
THIS RECORD ALL CITATIONS AVAILABLE IN THE RE  
FORMAT

L6 ANSWER 16 OF 63 CAPLUS COPYRIGHT 2006 ACS on STN  
AN 2002:552850 CAPLUS  
DN 137:213057

TI Silver-enhanced imaging of \*\*\*DNA\*\*\* hybridization at  
\*\*\*DNA\*\*\* microarrays with scanning electrochemical  
microscopy

AU Wang, Jun; Song, Fayi; Zhou, Feimeng

CS Department of Chemistry and Biochemistry, California State  
University, Los Angeles, CA, 90032, USA

SO Langmuir (2002), 18(17), 6653-6658 CODEN: LANGD5;  
ISSN: 0743-7463

PB American Chemical Society  
DT Journal  
LA English

AB The use of scanning electrochem. microscopy (SECM) to  
image oligodeoxynucleotide (ODN) hybridization at spots on  
microarrays has been demonstrated. ODN probes at a microarray  
surface were hybridized with a biotinylated target, and the  
regions where sequence-specific \*\*\*hybridization\*\*\* had  
occurred were developed by the silver staining process  
(adsorption of streptavidin-gold \*\*\*nanoparticles\*\*\* followed  
by silver particle deposition). As a consequence of the staining  
process, the surface cond. of the region where hybridization had  
taken place increased. Such an increase in cond. was sensitively  
detected by a SECM tip. The SECM detection level for a 17mer  
target was found to be at 30 amol per spot (or 3.0 fmol per  
slide). These values compare well with those from other  
detection methods (e.g., fluorescence and colorimetric  
detections). Coupled with the alteration of the hybridization  
temp., sequence-specific (single-base mismatch) \*\*\*DNA\*\*\*  
anal. can be accomplished. A reasonable sample throughput  
(imaging an area of 0.24 cm .times. 0.24 cm in about 38 min at a  
tip scanning speed of 50 .mu.m/s) was obtained.

RE.CNT 46 THERE ARE 46 CITED REFERENCES AVAILABLE  
FOR THIS RECORD ALL CITATIONS AVAILABLE IN THE RE  
FORMAT

L6 ANSWER 17 OF 63 CAPLUS COPYRIGHT 2006 ACS on STN  
AN 2002:479508 CAPLUS  
DN 137:197738

TI Sensitivity enhancement of \*\*\*DNA\*\*\* sensors by  
nanogold surface modification

AU Liu, Tao; Tang, Ji'an; Jiang, Long

CS Laboratory of Colloid and Interface Science, Chinese  
Academy of Science, Institute of Chemistry, Center for Molecular  
Science, Beijing, 100080, Peop. Rep. China

SO Biochemical and Biophysical Research Communications  
(2002), 295(1), 14-16 CODEN: BBRC9; ISSN: 0006-291X  
PB Elsevier Science

DT Journal  
LA English

AB A novel amplified microgravimetric gene sensing system was  
developed using quartz crystal microbalance modified by gold  
\*\*\*nanoparticles\*\*\* anchored on its 1,6-hexanedithiol modified  
gold electrode surface, and ultrasensitive detection of  
\*\*\*DNA\*\*\* \*\*\*hybridization\*\*\* was accomplished at the  
level of at least 2.times.10<sup>-16</sup> M.

RE.CNT 16 THERE ARE 16 CITED REFERENCES AVAILABLE  
FOR THIS RECORD ALL CITATIONS AVAILABLE IN THE RE  
FORMAT

L6 ANSWER 18 OF 63 CAPLUS COPYRIGHT 2006 ACS on STN  
AN 2002:429752 CAPLUS  
DN 137:75410

TI Particle size effect of the \*\*\*DNA\*\*\* sensor amplified  
with gold nanoparticles

AU Liu, Tao; Tang, Ji'an; Zhao, Hongqiu; Deng, Yongpei; Jiang,  
Long

CS Laboratory of Colloid and Interface Science, Center for  
Molecular Science, Institute of Chemistry, Chinese Academy of  
Science, Beijing, 100080, Peop. Rep. China

SO Langmuir (2002), 18(14), 5624-5626 CODEN: LANGD5;  
ISSN: 0743-7463

PB American Chemical Society  
DT Journal  
LA English

AB In this paper, nanoparticles with diams. ranging from 12 to 65 nm were used as amplifiers of a \*\*\*DNA\*\*\* sensor on a QCM device and the amplification efficiency was detd. Furthermore, gold \*\*\*nanoparticles\*\*\* with a certain size were immobilized on a modified mica slide by (3-mercaptopropyl)trimethoxysilane, and \*\*\*DNA\*\*\* hybridization was carried out on the surface. At. force microscopy (AFM) detection for the first time was used to intuitively observe the obvious surface changes at the mol. level during the hybridizing process.

RE.CNT 51 THERE ARE 51 CITED REFERENCES AVAILABLE FOR THIS RECORD ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 19 OF 63 CAPLUS COPYRIGHT 2006 ACS on STN  
AN 2002:409358 CAPLUS  
DN 137:89013

TI \*\*\*Hybridization\*\*\* and Enzymatic Extension of Au \*\*\*Nanoparticle\*\*\* -Bound Oligonucleotides

AU Pena, Sheila R. Nicewarner; Raina, Surabhi; Goodrich, Glenn P.; Fedoroff, Nina V.; Keating, Christine D.

CS Department of Chemistry and Life Sciences Consortium, The Pennsylvania State University, University Park, PA, 16802, USA  
SO Journal of the American Chemical Society (2002), 124(25), 7314-7323 CODEN: JACSAT; ISSN: 0002-7863

PB American Chemical Society  
DT Journal  
LA English

AB We have investigated the impact of steric effects on the hybridization and enzymic extension of oligonucleotides bound to 12-nm colloidal Au particles. In these expts., a \*\*\*nanoparticle\*\*\* -bound 12-mer sequence is \*\*\*hybridized\*\*\* either to its soln. phase 12-mer complement or to an 88-mer template sequence. The particle-bound oligonucleotide serves as a primer for enzymic extension reactions, in which covalent incorporation of nucleotides to form the complement of the template is achieved by the action of \*\*\*DNA\*\*\* polymerase. Primers were attached via-C6H12SH, -C12H24SH, and -TTACAATC6H12SH linkers attached at the 5' end. Primer coverage on the nanoparticles was varied by diln. with 5'HSC6H12AAA AAA3'. Hybridization efficiencies were detd. as a function of linker length, primer coverage, complement length (12-mer vs 88-mer), and primer:complement concn. ratio. In all cases, hybridization for the 88-mer was less efficient than for the 12-mer. Low primer surface coverages, greater particle-primer sepn., and higher primer:complement ratios led to optimal hybridization. Hybridization efficiencies as high as 98% and 75% were obsd. for the 12-mer and 88-mer, resp. Enzymic extension of particle-bound primers was obsd. under all conditions tested; however, the efficiency of the reaction was strongly affected by linker length and primer coverage. Extension of primers attached by the longest linker was as efficient as the soln.-phase reaction.  
RE.CNT 67 THERE ARE 67 CITED REFERENCES AVAILABLE FOR THIS RECORD ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 20 OF 63 CAPLUS COPYRIGHT 2006 ACS on STN  
AN 2002:405606 CAPLUS  
DN 138:148217

TI Recognition of \*\*\*DNA\*\*\* sequence and chain length by using \*\*\*DNA\*\*\* -linked nanoparticle  
AU Tang, Zhonglan; Mori, Takeshi; Takarada, Tohru; Maeda, Mizuo

CS Department of Applied Chemistry, Graduate School of Engineering, Kyushu University, Fukuoka, 812-8581, Japan  
SO Analytical Sciences (2001), 17(Suppl.), a357-a359 CODEN: ANSCEN; ISSN: 0910-6340

PB Japan Society for Analytical Chemistry  
DT Journal; (computer optical disk)  
LA English

AB We have studied the assembly of amphiphilic copolymer between oligonucleotide (ODN) as hydrophilic part and thermo-responsive polymer poly(N-isopropylacrylamide) (polyNIPAAm) as hydrophobic part. The copolymer formed colloidal nanoparticles, which aggregated depending on the sequence and chain length of coexisting \*\*\*DNA\*\*\*. The \*\*\*DNA\*\*\* -linked nanoparticle may be applied to an oligonucleotides discrimination system for gene diagnosis.

RE.CNT 8 THERE ARE 8 CITED REFERENCES AVAILABLE FOR THIS RECORD ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 21 OF 63 CAPLUS COPYRIGHT 2006 ACS on STN  
AN 2002:403338 CAPLUS  
DN 137:258035

TI An electrochemical \*\*\*DNA\*\*\* hybridization\*\*\* detection assay based on a silver \*\*\*nanoparticle\*\*\* label  
AU Cai, Hong; Xu, Ying; Zhu, Ningning; He, Pingang; Fang, Yuzhi

CS Department of Chemistry, East China Normal University, Shanghai, 200062, Peop. Rep. China  
SO Analyst (Cambridge, United Kingdom) (2002), 127(6), 803-808 CODEN: ANALAO; ISSN: 0003-2654

PB Royal Society of Chemistry  
DT Journal  
LA English  
AB A novel, sensitive electrochem. \*\*\*DNA\*\*\* hybridization\*\*\* detection assay, using silver \*\*\*nanoparticles\*\*\* as the oligonucleotide labeling tag, is described. The assay relies on the \*\*\*hybridization\*\*\* of the target \*\*\*DNA\*\*\* with the silver \*\*\*nanoparticle\*\*\* -oligonucleotide \*\*\*DNA\*\*\* probe, followed by the release of the silver metal atoms anchored on the hybrids by oxidative metal dissoln. and the indirect detn. of the solubilized AgI ions by anodic stripping voltammetry (ASV) at a carbon fiber ultramicroelectrode. The influence of the relevant exptl. variables, including the surface coverage of the target oligonucleotide, the duration of the silver dissoln. steps and the parameters of the electrochem. stripping measurement of the silver(i) ions, is examd. and optimized. The combination of the remarkable sensitivity of the stripping metal anal. at the microelectrode with the large no. of silver(i) ions released from each \*\*\*DNA\*\*\* hybrid allows detection at levels as low as 0.5 pmol L-1 of the target oligonucleotides.

RE.CNT 28 THERE ARE 28 CITED REFERENCES AVAILABLE FOR THIS RECORD ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 22 OF 63 CAPLUS COPYRIGHT 2006 ACS on STN  
AN 2002:392348 CAPLUS  
DN 137:164287

TI Sorting Fluorescent Nanocrystals with \*\*\*DNA\*\*\*

AU Gerion, Daniele; Parak, Wolfgang J.; Williams, Shara C.; Zanchet, Daniela; Micheel, Christine M.; Alivisatos, A. Paul  
CS Department of Chemistry, University of California, Berkeley, CA, 94720, USA

SO Journal of the American Chemical Society (2002), 124(24), 7070-7074 CODEN: JACSAT; ISSN: 0002-7863  
PB American Chemical Society



DT Journal  
LA English

AB Semiconductor nanocrystals with narrow and tunable fluorescence are covalently linked to oligonucleotides. These biocompounds retain the properties of both nanocrystals and \*\*\*DNA\*\*\*. Therefore, different sequences of \*\*\*DNA\*\*\* can be coded with nanocrystals and still preserve their ability to hybridize to their complements. We report the case where four different sequences of \*\*\*DNA\*\*\* are linked to four nanocrystal samples having different colors of emission in the range of 530-640 nm. When the \*\*\*DNA\*\*\*-nanocrystal conjugates are mixed together, it is possible to sort each type of \*\*\*nanoparticle\*\*\* by using \*\*\*hybridization\*\*\* on a defined micrometer-size surface contg. the complementary oligonucleotide. Detection of sorting requires only a single excitation source and an epifluorescence microscope. The possibility of directing fluorescent nanocrystals toward specific biol. targets and detecting them, combined with their superior photostability compared to org. dyes, opens the way to improved biolabeling expts., such as gene mapping on a nanometer scale or multicolor microarray anal.  
RE.CNT 24 THERE ARE 24 CITED REFERENCES AVAILABLE FOR THIS RECORD ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 23 OF 63 CAPLUS COPYRIGHT 2006 ACS on STN  
AN 2002:317810 CAPLUS  
DN 137:140724  
TI Multiple thiol-anchor capped \*\*\*DNA\*\*\*-gold nanoparticle conjugates  
AU Li, Zhi; Jin, Rongchao; Mirkin, Chad A.; Letsinger, Robert L.  
CS Department of Chemistry and Institute for Nanotechnology, Northwestern University, Evanston, IL, 60208, USA  
SO Nucleic Acids Research (2002), 30(7), 1558-1562 CODEN: NARHAD; ISSN: 0305-1048  
PB Oxford University Press  
DT Journal  
LA English  
OS CASREACT 137:140724  
AB We report the synthesis of a novel trithiol-capped oligodeoxyribonucleotide and gold nanoparticle conjugates prepd. from it. These \*\*\*DNA\*\*\*-gold \*\*\*nanoparticle\*\*\* conjugates exhibit substantially higher stability than analogs prepd. from monothiol and cyclic disulfide-capped oligodeoxyribonucleotides, but comparable \*\*\*hybridization\*\*\* properties. A quant. anal. of their stability under a range of conditions is provided. Significantly, this novel trithiol oligodeoxyribonucleotide can be used to stabilize particles >30 nm in diam., which are essential for many diagnostic applications.  
RE.CNT 24 THERE ARE 24 CITED REFERENCES AVAILABLE FOR THIS RECORD ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 24 OF 63 CAPLUS COPYRIGHT 2006 ACS on STN  
AN 2002:294945 CAPLUS  
DN 138:35445  
TI Gold nanoparticles as novel label for \*\*\*DNA\*\*\* diagnostics  
AU Csaki, Andrea; Moller, Robert; Fritzsche, Wolfgang  
CS Molecular Nanotechnology Group, Institute for Physical High Technology, Jena, 07702, Germany  
SO Expert Review of Molecular Diagnostics (2002), 2(2), 187-193 CODEN: ERMDCW; ISSN: 1473-7159  
PB Future Drugs Ltd.  
DT Journal; General Review  
LA English

AB A review. The growing interest in \*\*\*DNA\*\*\* diagnostics, esp. in combination with the need for highly-paralleled and miniaturized hybridization assays, is today addressed by fluorescence \*\*\*DNA\*\*\* chips. Fluorescence detection is approved and highly developed, however, it has also problematic aspects, e.g., the low stability of the dyes, the influence of the physicochem. environment onto the signal intensity and the expensive set-up for detection. A novel detection scheme based on metal nanoparticles was proposed to overcome these problems and is discussed in this review.  
RE.CNT 27 THERE ARE 27 CITED REFERENCES AVAILABLE FOR THIS RECORD ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 25 OF 63 CAPLUS COPYRIGHT 2006 ACS on STN  
AN 2002:293191 CAPLUS  
DN 137:289398  
TI Gene diagnosis and nanotechnology  
AU Tang, Zhonglan; Maeda, Mizuo  
CS Graduate School of Engineering, Kyushu University, Japan  
SO Bio Industry (2002), 19(3), 16-22 CODEN: BIINEG; ISSN: 0910-6545  
PB Shi Emu Shi Shuppan  
DT Journal; General Review  
LA Japanese

AB A review. The SNP anal. by \*\*\*DNA\*\*\* \*\*\*hybridization\*\*\* using the allele-specific \*\*\*DNA\*\*\* probe immobilized on gold \*\*\*nanoparticles\*\*\* was described. Spectrophotometric detection of \*\*\*nanoparticle\*\*\* aggregation upon \*\*\*DNA\*\*\* \*\*\*hybridization\*\*\* and SNP detection based on the difference in phase transfer process of the colloidal particles between the matched and the mismatched \*\*\*DNA\*\*\* pair combination were described. Detailed examn. of the system set-ups (optimum particle concn., temp., salt concn., etc.) that had been conducted for sensing slight difference in m.ps. between wild and mutant types was presented. Sensitive detection system using probe-immobilized cantilever sensors to detect phys. stress upon hybridization was also described as another example of nanotechnol. approach.

L6 ANSWER 26 OF 63 CAPLUS COPYRIGHT 2006 ACS on STN  
AN 2002:234786 CAPLUS  
DN 136:242425  
TI Magnetically-induced solid-state electrochemical detection of \*\*\*DNA\*\*\* hybridization  
AU Wang, Joseph; Xu, Danke; Polsky, Ronen  
CS Department of Chemistry and Biochemistry, New Mexico State University, Las Cruces, NM, 88003, USA  
SO Journal of the American Chemical Society (2002), 124(16), 4208-4209 CODEN: JACSAT; ISSN: 0002-7863  
PB American Chemical Society  
DT Journal  
LA English

AB A magnetic triggering of a solid-state elec. transduction of \*\*\*DNA\*\*\* hybridization is described. Positioning of an external magnet below the thick-film electrode attracts the \*\*\*DNA\*\*\* /particle network and enables the solid-state electrochem. stripping detection of the silver tracer. TEM imaging indicates that the \*\*\*hybridization\*\*\* event results in a three-dimensional aggregate structure in which duplex segments link the metal \*\*\*nanoparticles\*\*\* and magnetic spheres, and that most of this assembly is covered with the silver ppt. This leads to a direct contact of the metal tag with the surface (in connection to the magnetic collection) and enables the solid-state electrochem. transduction (without prior dissoln. and subsequent electrodeposition of the metal), using oxidative



dissohn. of the silver tracer. No such aggregates (and hence magnetic "collection") are obsd. in the presence of noncomplementary \*\*\*DNA\*\*\*, i.e., without the linking hybrid. The new method couples high sensitivity of silver-amplified assays with effective discrimination against excess of closely related nucleotide sequences (including single-base imperfections). Such direct elec. detection of \*\*\*DNA\*\*\*/metal-particle assemblies can bring new capabilities to the detection of \*\*\*DNA\*\*\* hybridization, and could be applied to other bioaffinity assays.

RE.CNT 13 THERE ARE 13 CITED REFERENCES AVAILABLE FOR THIS RECORD ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 27 OF 63 CAPLUS COPYRIGHT 2006 ACS on STN AN 2002:212207 CAPLUS DN 136:382462

TI Bio-barcodes based on oligonucleotide-modified nanoparticles

AU Nam, Jwa-Min; Park, So-Jung; Mirkin, Chad A.  
CS Department of Chemistry and Institute for Nanotechnology, Northwestern University, Evanston, IL, 60208, USA  
SO Journal of the American Chemical Society (2002), 124(15), 3820-3821 CODEN: JACSAT; ISSN: 0002-7863

PB American Chemical Society

DT Journal

LA English

AB By utilizing oligonucleotide-modified Au nanoparticles encoded with sequences that act as biobarcode, one can screen for multiple target polyvalent proteins simultaneously in one soln. This novel concept was demonstrated with two types of detection formats, a homogeneous assay and one based on oligonucleotide microarrays. With such an approach, one can prep. an extraordinarily large no. of barcodes from synthetically accessible oligonucleotides (e.g., a 12-mer sequence offers 412 possible barcodes).

RE.CNT 14 THERE ARE 14 CITED REFERENCES AVAILABLE FOR THIS RECORD ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 28 OF 63 CAPLUS COPYRIGHT 2006 ACS on STN AN 2002:188392 CAPLUS

TI Surface effect control: improve the \*\*\*hybridization\*\*\* efficiency of the immobilized molecular beacon and other \*\*\*DNA\*\*\* probes on \*\*\*nanoparticles\*\*\*

AU Li, Jianwei J.; Tan, Weihong  
CS Department of Chemistry, University of Florida, Gainesville, FL, 32611, USA

SO Abstracts of Papers, 223rd ACS National Meeting, Orlando, FL, United States, April 7-11, 2002 (2002), COLL-474 Publisher: American Chemical Society, Washington, D. C. CODEN: 69CKQP

DT Conference; Meeting Abstract

LA English

AB There is growing interest in the exploitation of high-throughput solid-phase-based assays for diagnosis, new drugs development and biomedical research. Techniques using solid support-bound oligonucleotides as probes are finding a wide range of applications. Of all of these applications, hybridization of tethered oligonucleotide probes with nucleic acid targets is the central event, and high hybridization efficiency holds the key for sensitive, reproducible and reliable assays. Whatever solid supports is chosen, a common problem to be faced is that the solid support always tends to constrain the ways in which the bound oligonucleotide probes can interact with target mols. Effort has been made to mitigate the influence of solid support. In this paper, we have developed a novel approach to control surface

effect to improve \*\*\*DNA\*\*\* \*\*\*hybridization\*\*\* efficiency on \*\*\*nanoparticle\*\*\* surfaces. This approach significantly increases the hybridization speed and yield, and should be generally useful for all solid-phase-based \*\*\*DNA\*\*\* assays.

L6 ANSWER 29 OF 63 CAPLUS COPYRIGHT 2006 ACS on STN AN 2002:158929 CAPLUS DN 137:58213

TI Array-based electrical detection of \*\*\*DNA\*\*\* with nanoparticle probes

AU Park, So-Jung; Taton, T. Andrew; Mirkint, Chad A.  
CS Department of Chemistry and Institute for Nanotechnology, Northwestern University, Evanston, IL, 60208, USA  
SO Science (Washington, DC, United States) (2002), 295(5559), 1503-1506 CODEN: SCIEAS; ISSN: 0036-8075

PB American Association for the Advancement of Science

DT Journal

LA English

AB A \*\*\*DNA\*\*\* array detection method is reported in which the binding of oligonucleotides functionalized with gold nanoparticles leads to cond. changes assocd. with target-probe binding events. The binding events localize gold nanoparticles in an electrode gap; silver deposition facilitated by these nanoparticles bridges the gap and leads to readily measurable cond. changes. An unusual salt concn.-dependent \*\*\*hybridization\*\*\* behavior assocd. with these \*\*\*nanoparticle\*\*\* probes was exploited to achieve selectivity without a thermal-stringency wash. Using this method, we have detected target \*\*\*DNA\*\*\* at concns. as low as 500 femtomolar with a point mutation selectivity factor of .apprx.100,000:1.

RE.CNT 20 THERE ARE 20 CITED REFERENCES AVAILABLE FOR THIS RECORD ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 30 OF 63 CAPLUS COPYRIGHT 2006 ACS on STN AN 2002:151845 CAPLUS DN 136:275546

TI \*\*\*DNA\*\*\*-based magnetic nanoparticle assembly acts as a magnetic relaxation nanoswitch allowing screening of \*\*\*DNA\*\*\*-cleaving agents

AU Perez, J. Manuel; O'Loughin, Terence; Simeone, F. Joseph; Weissleder, Ralph; Josephson, Lee

CS MGH-Center for Molecular Imaging Research, Harvard Medical School, Charlestown, MA, 02129, USA

SO Journal of the American Chemical Society (2002), 124(12), 2856-2857 CODEN: JACSAT; ISSN: 0002-7863

PB American Chemical Society

DT Journal

LA English

AB Monodisperse magnetic nanoparticles conjugated with complementary oligonucleotide sequences self-assemble into stable magnetic nanoassemblies resulting in a decrease of the spin-spin relaxation times (T2) of neighboring water protons. When these nanoassemblies are treated with a \*\*\*DNA\*\*\* cleaving agent, the nanoparticles become dispersed, switching the T2 of the soln. back to original values. These qualities render the developed nanoparticles and their nanoassemblies as magnetic relaxation switches capable of screening for \*\*\*DNA\*\*\*-cleaving compds. by magnetic resonance methods such as MRI and NMR.

L6 ANSWER 31 OF 63 CAPLUS COPYRIGHT 2006 ACS on STN AN 2002:106225 CAPLUS DN 136:259512

TI \*\*\*Nanoparticle\*\*\* Layers Assembled through  
\*\*\*DNA\*\*\* \*\*\*Hybridization\*\*\* : Characterization and  
Optimization  
AU Sauthier, Marc L.; Carroll, R. Lloyd; Gorman, Christopher B.;  
Franzen, Stefan  
CS Department of Chemistry, North Carolina State University,  
Raleigh, NC, 27695, USA  
SO Langmuir (2002), 18(5), 1825-1830 CODEN: LANGD5; ISSN:  
0743-7463  
PB American Chemical Society  
DT Journal  
LA English  
AB The \*\*\*hybridization\*\*\* of \*\*\*nanoparticle\*\*\* -  
labeled \*\*\*DNA\*\*\* targets to surface-attached \*\*\*DNA\*\*\*  
probes has been investigated. Scanning tunneling microscopy  
(STM) and Raman and Fourier transform IR (FTIR) spectroscopy  
were used to elucidate surface morphol., coverage, and the  
presence of aggregates. The factors that affect surface  
coverage, such as probe d., labeled target concn., and particle  
size, were systematically investigated by STM in order to det.  
the best set of exptl. conditions allowing the formation of dense  
monolayers with a minimal no. of surface defects for both 5(+-.  
.1) nm and 10(+-.2) nm gold \*\*\*nanoparticle\*\*\* labels on  
the target strand. Grazing-angle FTIR spectroscopy demonstrates  
that \*\*\*DNA\*\*\* is largely oriented once the labeled targets  
\*\*\*hybridized\*\*\* to the probes. Raman microscopy was used  
to probe the surface for the presence of large aggregates that  
would give rise to large scattering signals. Both STM and optical  
expts. provide evidence that dense surface layers can be formed  
without extensive aggregation. Nonselective binding was shown  
to be a function of the target concn. and nanoparticle size.  
Propensity for both aggregation and nonspecific binding is  
greater for 10(+-.2) nm than for 5(+-.1) nm gold nanoparticles.  
RE.CNT 17 THERE ARE 17 CITED REFERENCES AVAILABLE  
FOR THIS RECORD ALL CITATIONS AVAILABLE IN THE RE  
FORMAT

L6 ANSWER 32 OF 63 CAPLUS COPYRIGHT 2006 ACS on STN  
AN 2001:922058 CAPLUS  
DN 136:131164  
TI Immobilization of gold \*\*\*nanoparticles\*\*\* on solid  
supports utilizing \*\*\*DNA\*\*\* \*\*\*hybridization\*\*\*  
AU Peschel, S.; Ceyhan, B.; Niemeyer, C. M.; Gao, S.; Chi, L.;  
Simon, U.  
CS Institut für Anorganische Chemie, RWTH Aachen, Aachen,  
52056, Germany  
SO Materials Science & Engineering, C: Biomimetic and  
Supramolecular Systems (2002), C19(1-2), 47-50 CODEN:  
MSCEEE; ISSN: 0928-4931  
PB Elsevier Science B.V.  
DT Journal  
LA English  
AB Self-organization of colloidal metal nanoparticles into micro-  
and nanostructured assemblies is currently of tremendous  
interest promising to find new size- and structure-dependent  
phys. properties. Owing to its unique recognition capabilities and  
physicochem. stability, \*\*\*DNA\*\*\* can be used as a mol.  
linker for gold nanoparticles and is a promising construction  
material for their precise spatial positioning. Due to the  
enormous specificity of nucleic acid hybridization, the site-specific  
immobilization of \*\*\*DNA\*\*\* -functionalized gold colloids (1-  
40 nm) to solid supports, previously functionalized with a  
complementary \*\*\*DNA\*\*\* array, allows the fabrication of  
novel nanostructured surface architectures. Scanning force  
microscopy (SFM), used to characterize the intermediate steps of  
the \*\*\*DNA\*\*\* -directed immobilization (DDI) on a gold

substrate, provides initial insight into the specificity and efficiency  
of this technique.  
RE.CNT 16 THERE ARE 16 CITED REFERENCES AVAILABLE  
FOR THIS RECORD ALL CITATIONS AVAILABLE IN THE RE  
FORMAT

L6 ANSWER 33 OF 63 CAPLUS COPYRIGHT 2006 ACS on STN  
AN 2001:837901 CAPLUS  
DN 136:65781  
TI The percolation transition in the \*\*\*DNA\*\*\* -gold  
nanoparticle system  
AU Kiang, Ching-Hwa; Ramos, Rona  
CS Dep. Physics and Astronomy, Univ. California, Los Angeles,  
CA, 90095-1547, USA  
SO Los Alamos National Laboratory, Preprint Archive, Physics  
(2001) 1-4, arXiv:physics/0111002, 1 Nov 2001 CODEN: LNPFP9  
URL: <http://xxx.lanl.gov/pdf/physics/0111002>  
PB Los Alamos National Laboratory  
DT Preprint  
LA English  
AB Melting and \*\*\*hybridization\*\*\* of \*\*\*DNA\*\*\* -  
capped gold \*\*\*nanoparticle\*\*\* networks are investigated  
with optical absorption spectroscopy and TEM. Single-stranded,  
12-base \*\*\*DNA\*\*\* -capped gold nanoparticles are linked with  
complementary, single-stranded, 24-base linker \*\*\*DNA\*\*\* to  
form particle networks. Compared to free \*\*\*DNA\*\*\*, a  
sharp melting transition is seen in these networked \*\*\*DNA\*\*\*  
-nanoparticle systems. The sharpness is explained by percolation  
transition phenomena.  
RE.CNT 22 THERE ARE 22 CITED REFERENCES AVAILABLE  
FOR THIS RECORD ALL CITATIONS AVAILABLE IN THE RE  
FORMAT

L6 ANSWER 34 OF 63 CAPLUS COPYRIGHT 2006 ACS on STN  
AN 2001:783614 CAPLUS  
DN 136:115014  
TI Formation of \*\*\*DNA\*\*\* -carrying colloidal particle from  
poly(N-isopropylacrylamide)-graft- \*\*\*DNA\*\*\* copolymer and  
its assembly through hybridization  
AU Mori, Takeshi; Maeda, Mizuo  
CS Department of Applied Chemistry, Graduate School of  
Engineering, Kyushu University, Fukuoka, 812-8581, Japan  
SO Polymer Journal (Tokyo, Japan) (2001), 33(10), 830-833  
CODEN: POLJBB; ISSN: 0032-3896  
PB Society of Polymer Science, Japan  
DT Journal  
LA English  
AB A one-step prepn. of \*\*\*DNA\*\*\* -carrying colloidal  
nanoparticle through the self-organization of copolymer,  
composed of poly(N- isopropylacrylamide) (PNIPAAm) main  
chain and \*\*\*DNA\*\*\* graft chain, is described. The narrowly  
distributed \*\*\*DNA\*\*\* -carrying colloidal particles are easily  
prepd. from PNIPAAm-graft- \*\*\*DNA\*\*\* by heating. The  
particle surface \*\*\*DNA\*\*\* recognizes the complementary  
crosslinking \*\*\*DNA\*\*\* so that the particle assembly is  
formed. This particle would be applicable for the turbidimetric  
\*\*\*DNA\*\*\* detection.  
RE.CNT 22 THERE ARE 22 CITED REFERENCES AVAILABLE  
FOR THIS RECORD ALL CITATIONS AVAILABLE IN THE RE  
FORMAT

L6 ANSWER 35 OF 63 CAPLUS COPYRIGHT 2006 ACS on STN  
AN 2001:745088 CAPLUS  
DN 135:267870  
TI Metal \*\*\*nanoparticle\*\*\* -based electrochemical stripping  
potentiometric detection of \*\*\*DNA\*\*\* \*\*\*hybridization\*\*\*

AU Wang, Joseph; Xu, Danke; Kawde, Abdel-Nasser; Polsky, Ronen  
CS Department of Chemistry and Biochemistry, New Mexico State University, Las Cruces, NM, 88003, USA  
SO Analytical Chemistry (2001), 73(22), 5576-5581 CODEN: ANCHAM; ISSN: 0003-2700  
PB American Chemical Society  
DT Journal  
LA English  
AB A new \*\*\*nanoparticle\*\*\* -based elec. detection of \*\*\*DNA\*\*\* \*\*\*hybridization\*\*\*, based on electrochem. stripping detection of the colloidal gold tag, is described. In this protocol, the \*\*\*hybridization\*\*\* of a target oligonucleotide to magnetic bead-linked oligonucleotide probes is followed by binding of the streptavidin-coated metal \*\*\*nanoparticles\*\*\* to the captured \*\*\*DNA\*\*\*, dissoln. of the nanometer-sized gold tag, and potentiometric stripping measurements of the dissolved metal tag at single-use thick-film carbon electrodes. An advanced magnetic processing technique is used to isolate the \*\*\*DNA\*\*\* duplex and to provide low-vol. mixing. The influence of relevant exptl. variables, including the amts. of the gold \*\*\*nanoparticles\*\*\* and the magnetic beads, the duration of the \*\*\*hybridization\*\*\* and gold dissoln. steps, and the parameters of the potentiometric stripping operation upon the hybridization signal, is examd. and optimized. Transmission electron microscopy micrographs indicate that the \*\*\*hybridization\*\*\* event leads to the bridging of the gold \*\*\*nanoparticles\*\*\* to the magnetic beads. Further signal amplification, and lowering of the detection limits to the nanomolar and picomolar domains, are achieved by pptg. gold or silver, resp., onto the colloidal gold label. The new electrochem. stripping metallogenomagnetic protocol couples the inherent signal amplification of stripping metal anal. with discrimination against nonhybridized \*\*\*DNA\*\*\*, the use of microliter sample vols., and disposable transducers and, hence, offers great promise for decentralized genetic testing.  
RE.CNT 30 THERE ARE 30 CITED REFERENCES AVAILABLE FOR THIS RECORD ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 36 OF 63 CAPLUS COPYRIGHT 2006 ACS on STN AN 2001:685101 CAPLUS  
DN 136:319894  
TI \*\*\*Nanoparticle\*\*\* -amplified surface plasmon resonance for detection of \*\*\*DNA\*\*\* \*\*\*hybridization\*\*\*  
AU Goodrich, Glenn P.; Nicewarner, Sheila R.; He, Lin; Natan, Michael J.; Keating, Christine D.  
CS The Pennsylvania State University, University Park, PA, 16802, USA  
SO Proceedings of SPIE-The International Society for Optical Engineering (2001), 4258(Nanoparticles and Nanostructured Surfaces: Novel Reporters with Biological Applications), 80-85 CODEN: PSISDG; ISSN: 0277-786X  
PB SPIE-The International Society for Optical Engineering  
DT Journal  
LA English  
AB In recent years there has been a great deal of interest in the measurement of \*\*\*DNA\*\*\* hybridization at surfaces. Surface-confined \*\*\*DNA\*\*\* hybridization has been used to monitor gene expression, to detect the presence of a particular \*\*\*DNA\*\*\* sequence and det. single nucleotide polymorphisms (SNPs). \*\*\*DNA\*\*\* microarrays, which can contain thousands of discrete \*\*\*DNA\*\*\* sequences on a single surface, have become widely used for hybridization studies. While a powerful technique, this technol. is limited by the stability of the fluorescent dyes used to label the \*\*\*DNA\*\*\*, and the

need to perform measurements ex-situ to reduce the fluorescence background. In this report, we describe the use of colloid-amplified surface plasmon resonance (SPR) to measure \*\*\*DNA\*\*\* hybridization at surfaces. SPR is a surface sensitive technique, which can be used to study hybridization in situ, and the use of colloidal metal tags provides excellent sensitivity. Angle-scanning SPR has been used to study oligonucleotide hybridization to surface confined probes, and work is underway to apply SPR imaging to study \*\*\*DNA\*\*\* hybridization in macro- and microarray formats.  
RE.CNT 26 THERE ARE 26 CITED REFERENCES AVAILABLE FOR THIS RECORD ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 37 OF 63 CAPLUS COPYRIGHT 2006 ACS on STN AN 2001:676892 CAPLUS  
DN 135:237555  
TI Integrated nucleic acid hybridization devices for improved kinetics, sensitivity and discrimination power  
IN Hogan, Michael; Powdrill, Thomas; Iverson, Bonnie; Belosludtsev, Yuri Y.; Belosludtsev, Inna Y.  
PA Genometrix Genomix, Inc., USA  
SO PCT Int. Appl., 101 pp. CODEN: PIXXD2  
DT Patent  
LA English  
FAN.CNT 1 PATENT NO. KIND DATE APPLICATION  
NO. DATE -----  
PI WO 2001066687 A1 20010913 WO 2000-US23438  
20000824 W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG  
PRAI US 2000-522240 A1 20000309 US 2000-636268  
A1 20000810  
AB The invention provides devices and methods for enhanced and selective assocn. or binding between biol. materials, such as nucleic acids, e.g., \*\*\*DNA\*\*\* or \*\*\*RNA\*\*\*, or polypeptides, and an immobilized oligonucleotide probe. In one embodiment, the invention provides an assocn. device comprising a plurality of nucleic acid probes or polypeptide probes or a combination thereof linked to a solid substrate. The solid substrate comprises a substrate surface comprising an external substrate surface and a plurality of internal pores, wherein the pores comprise a proximal end opening to the external surface to allow passage of fluid into a pore, and wherein the pore surfaces comprise an assocn. surface. The assocn. surface comprises a charged surface comprising net pos. (cationic) charge d. under conditions comprising a pH lower than the pI of the assocn. surface. Methods for making these hybridization/assocn. devices are also provided. Covalent and noncovalent probe immobilization methodologies are employed for surface hybridization modeling studies. Incorporating low ionic strength, low pH buffers (together with a net cationic charge d. on the device surface) as hybridization conditions provides significant increases in the kinetics, sensitivity, and discrimination power of nucleic acid-based and polypeptide-based biosensors and related hybridization techniques. For example, the devices and methods of the invention can be used in nucleic acid-based diagnostic tests. The devices and methods of the invention can be used,

e.g., for detecting the assocn. of a nucleic acid in a sample to a nucleic acid probe or purifying a nucleic acid from a sample.  
RE.CNT 6 THERE ARE 6 CITED REFERENCES AVAILABLE FOR THIS RECORD ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 38 OF 63 CAPLUS COPYRIGHT 2006 ACS on STN  
AN 2001:638227 CAPLUS  
TI Oligonucleotide-nanoparticle conjugates with multiple anchor groups  
AU Li, Zhi; Jin, Rongchao; Mirkin, Chad A.; Letsinger, Robert L.  
CS Chemistry Department and Center for Nanofabrication and Molecular Self-assembly, Northwestern University, Evanston, IL, 60208, USA  
SO Abstracts of Papers, 222nd ACS National Meeting, Chicago, IL, United States, August 26-30, 2001 (2001), COLL-191  
Publisher: American Chemical Society, Washington, D. C. CODEN: 69BUZP  
DT Conference; Meeting Abstract  
LA English  
AB Conjugates of oligonucleotides and nanoparticles are gaining importance in both material science and biotechnol. as novel building blocks and sensitive diagnostic probes. Diagnostic applications of such conjugates in particular demand high stability in media contg. strong surface-binding agents, such as alkanethiols. Conventional alkanethiol-capped \*\*\*DNA\*\*\* - modified gold nanoparticles show considerable irreversible aggregation in the presence of competing thiols due to loss of \*\*\*DNA\*\*\* from the surface. Herein we present alternative oligonucleotide attachment strategies based on a cyclic steroid disulfide and branched tri-hexylthiol modification of the \*\*\*DNA\*\*\* terminus. These capping groups afford substantial enhancement of particle stability over single hexanethiol anchors. We will compare the stability, surface modification properties, and \*\*\*hybridization\*\*\* activity of gold \*\*\*nanoparticle\*\*\* probes with these novel surface modifications.

L6 ANSWER 39 OF 63 CAPLUS COPYRIGHT 2006 ACS on STN  
AN 2001:621648 CAPLUS  
DN 135:371949  
TI Directed assembly of periodic materials from protein and oligonucleotide-modified nanoparticle building blocks  
AU Park, So-Jung; Lazarides, Anne A.; Mirkin, Chad A.; Letsinger, Robert L.  
CS Department of Chemistry and Center for Nanofabrication and Molecular Self Assembly, Northwestern University, Evanston, IL, 60208-3113, USA  
SO Angewandte Chemie, International Edition (2001), 40(15), 2909-2912 CODEN: ACIEF5; ISSN: 1433-7851  
PB Wiley-VCH Verlag GmbH  
DT Journal  
LA English  
AB \*\*\*DNA\*\*\* -directed assembly of \*\*\*nanoparticles\*\*\* was achieved by linking thio-alkyl-substituted oligodeoxynucleotide chains to gold \*\*\*nanoparticles\*\*\* or biotin-substituted oligodeoxynucleotides to streptavidin, and then \*\*\*hybridizing\*\*\* the two with a complimentary oligodeoxynucleotide linker. The thermal dissocn. of the aggregates showed features of both aggregate particle growth and \*\*\*DNA\*\*\* melting; one method of increasing the size of aggregates formed was to heat the mixt. to a few degrees below the m.p.

L6 ANSWER 40 OF 63 CAPLUS COPYRIGHT 2006 ACS on STN  
AN 2001:610135 CAPLUS  
DN 135:190978

TI Silver-Enhanced Colloidal Gold Electrochemical Stripping Detection of \*\*\*DNA\*\*\* Hybridization  
AU Wang, Joseph; Polsky, Ronen; Xu, Danke  
CS Department of Chemistry and Biochemistry, New Mexico State University, Las Cruces, NM, 88003, USA  
SO Langmuir (2001), 17(19), 5739-5741 CODEN: LANGD5; ISSN: 0743-7463  
PB American Chemical Society  
DT Journal  
LA English  
AB We report on a novel method for detecting \*\*\*DNA\*\*\* \*\*\*hybridization\*\*\*, based on the pptn. of silver on gold \*\*\*nanoparticle\*\*\* tags and a subsequent electrochem. stripping detection of the dissolved silver. Such coupling of a \*\*\*nanoparticle\*\*\* -promoted silver pptn. with the remarkable sensitivity of stripping metal anal. offers a dramatic enhancement of the \*\*\*hybridization\*\*\* response. An efficient magnetic isolation of the duplex is used for discriminating against nonhybridized \*\*\*DNA\*\*\*, including an excess of mismatched oligonucleotides. The new silver-enhanced colloidal gold stripping detection strategy holds great promise for the detection of \*\*\*DNA\*\*\* hybridization and represents an attractive alternative to indirect optical affinity assays of nucleic acids and other biomols.  
RE.CNT 12 THERE ARE 12 CITED REFERENCES AVAILABLE FOR THIS RECORD ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 41 OF 63 CAPLUS COPYRIGHT 2006 ACS on STN  
AN 2001:578956 CAPLUS  
DN 135:164328  
TI Gold nanoparticle-based quantitative electrochemical detection of amplified human cytomegalovirus \*\*\*DNA\*\*\* using disposable microband electrodes  
AU Authier, Laurent; Grossiord, Celine; Brossier, Pierre; Limoges, Benoit  
CS Laboratoire de Microbiologie Medicale et Moleculaire Faculte de Medecine et de Pharmacie, Faculte de Medecine et de Pharmacie, Dijon, 21033, Fr.  
SO Analytical Chemistry (2001), 73(18), 4450-4456 CODEN: ANCHAM; ISSN: 0003-2700  
PB American Chemical Society  
DT Journal  
LA English  
AB An electrochem. \*\*\*DNA\*\*\* detection method has been developed for the sensitive quantification of an amplified 406-base pair human cytomegalovirus \*\*\*DNA\*\*\* sequence (HCMV \*\*\*DNA\*\*\*). The assay relies on (i) the \*\*\*hybridization\*\*\* of the single-stranded target HCMV \*\*\*DNA\*\*\* with an oligonucleotide-modified Au \*\*\*nanoparticle\*\*\* probe, (ii) followed by the release of the gold metal atoms anchored on the hybrids by oxidative metal dissoln., and (iii) the indirect detn. of the solubilized AuIII ions by anodic stripping voltammetry at a sandwich-type screen-printed microband electrode (SPMBE). Due to the enhancement of the AuIII mass transfer by nonlinear diffusion during the electrodeposition time, the SPMBE allows the sensitive detn. of AuIII in a small vol. of quiescent soln. The combination of the sensitive AuIII detn. at a SPMBE with the large no. of AuIII released from each gold nanoparticle probe allows detection of as low as 5 pM amplified HCMV \*\*\*DNA\*\*\* fragment.  
RE.CNT 45 THERE ARE 45 CITED REFERENCES AVAILABLE FOR THIS RECORD ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 42 OF 63 CAPLUS COPYRIGHT 2006 ACS on STN

AN 2001:530445 CAPLUS  
DN 135:207798  
TI \*\*\*DNA\*\*\* -modified core-shell Ag/Au nanoparticles  
AU Cao, YunWei; Jin, Rongchao; Mirkin, Chad A.  
CS Department of Chemistry and Center for Nanofabrication and Molecular Self-assembly, Northwestern University, Evanston, IL, 60208, USA  
SO Journal of the American Chemical Society (2001), 123(32), 7961-7962 CODEN: JACSAT; ISSN: 0002-7863  
PB American Chemical Society  
DT Journal  
LA English  
AB In this article the authors report a low temp. method for generating core-shell particles consisting of a core of Ag and a monolayer shell of Au that can be readily functionalized with oligonucleotides using the proven preparatory methods for pure gold particle oligonucleotide conjugates. Moreover, the authors show how this novel nanoparticle compn. can be used to access a colorimetric detections system distinct from pure gold systems.  
RE.CNT 22 THERE ARE 22 CITED REFERENCES AVAILABLE FOR THIS RECORD ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 43 OF 63 CAPLUS COPYRIGHT 2006 ACS on STN  
AN 2001:424278 CAPLUS  
DN 135:233111  
TI Scanning electron microscope observation of heterogeneous three-dimensional nanoparticle arrays using \*\*\*DNA\*\*\*  
AU Takahagi, Takayuki; Tsutsui, Gen; Huang, Shujuan; Sakaue, Hiroyuki; Shingubara, Shoso  
CS Graduate School of Advanced Sciences of Matter, Hiroshima University, Higashi-Hiroshima, 739-8526, Japan  
SO Japanese Journal of Applied Physics, Part 2: Letters (2001), 40(5B), L521-L523 CODEN: JAPLD8; ISSN: 0021-4922  
PB Japan Society of Applied Physics  
DT Journal  
LA English  
AB \*\*\*DNA\*\*\* oligonucleotides are considered to be as a useful tool for fabricating complex structures on a nanometer scale because of their selective reactivity. Herein, the authors describe a fabrication technique for heterogeneous three-dimensional nanoparticle arrays composed of Au nanoparticles of two different sizes linked by thiol-synthesized \*\*\*DNA\*\*\* oligonucleotides. Each size of the Au nanoparticles was encapsulated by complementary \*\*\*DNA\*\*\* oligonucleotides. \*\*\*DNA\*\*\* hybridization occurs upon their mixing, resulting in the construction of a three-dimensional nanostructure. Scanning electron microscope (SEM) observation and UV spectroscopy were performed to confirm the construction of the nanostructure. This fabrication technique will be crucial for the advancement of nanotechnol. in the next ten years.  
RE.CNT 30 THERE ARE 30 CITED REFERENCES AVAILABLE FOR THIS RECORD ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 44 OF 63 CAPLUS COPYRIGHT 2006 ACS on STN  
AN 2001:320365 CAPLUS  
DN 134:321368  
TI Two-Color Labeling of Oligonucleotide Arrays via Size-Selective Scattering of Nanoparticle Probes  
AU Taton, T. Andrew; Lu, Gang; Mirkin, Chad A.  
CS Department of Chemistry and Center for Nanofabrication and Molecular Self-Assembly, Northwestern University, Evanston, IL, 60208, USA

SO Journal of the American Chemical Society (2001), 123(21), 5164-5165 CODEN: JACSAT; ISSN: 0002-7863  
PB American Chemical Society  
DT Journal  
LA English  
AB Imaging the light scattered by oligonucleotide-functionalized, 50 and 100 nm diam. gold \*\*\*nanoparticle\*\*\* probes \*\*\*hybridized\*\*\* to targets captured by a \*\*\*DNA\*\*\* array is reported. The method can be used to identify two target sequences in one soln. It is also shown that with the scanometric array method, the unique melting properties of nanometric probes lead to enhanced sequence selectivity when \*\*\*DNA\*\*\* arrays are imaged by scattering light from bound nanoparticles.  
RE.CNT 20 THERE ARE 20 CITED REFERENCES AVAILABLE FOR THIS RECORD ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 45 OF 63 CAPLUS COPYRIGHT 2006 ACS on STN  
AN 2001:228137 CAPLUS  
DN 135:327869  
TI A three-dimensional heterogeneous \*\*\*DNA\*\*\* sensing surface formed by attaching oligodeoxynucleotide-capped gold nanoparticles onto a gold-coated quartz crystal  
AU Han, Shubo; Lin, Jianqiao; Satjapipat, Munlika; Baca, Alfred J.; Zhou, Feimeng  
CS Department of Chemistry and Biochemistry, California State University, Los Angeles, Los Angeles, CA, 90032, USA  
SO Chemical Communications (Cambridge, United Kingdom) (2001), (7), 609-610 CODEN: CHCOFS; ISSN: 1359-7345  
PB Royal Society of Chemistry  
DT Journal  
LA English  
AB Exposing oligodeoxynucleotide (ODN)-capped Au nanoparticles to a quartz crystal under shear oscillation resulted in the formation of a uniform monolayer contg. these \*\*\*nanoparticles\*\*\* or multilayers with islands of the ODN-capped \*\*\*nanoparticles\*\*\*, which, in turn, improved the extent of \*\*\*DNA\*\*\* \*\*\*hybridization\*\*\*.  
RE.CNT 16 THERE ARE 16 CITED REFERENCES AVAILABLE FOR THIS RECORD ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 46 OF 63 CAPLUS COPYRIGHT 2006 ACS on STN  
AN 2001:199683 CAPLUS  
TI Nanoparticle-based bioanalysis  
AU Keating, Christine D.; He, Lin; Nicewarner, Sheila; Reiss, Brian D.; Pena, Dave; Goodrich, Glenn P.; Natan, Michael J.; Mallouk, Thomas E.  
CS Department of Chemistry, Pennsylvania State University, University Park, PA, 16802, USA  
SO Abstracts of Papers, 221st ACS National Meeting, San Diego, CA, United States, April 1-5, 2001 (2001) COLL-425 CODEN: 69FZD4  
PB American Chemical Society  
DT Journal; Meeting Abstract  
LA English  
AB Two applications of metal nanoparticles in biosensing are described. In the first, \*\*\*DNA\*\*\* hybridization at an Au surface is detected via colloidal Au amplified surface plasmon resonance (SPR). Use of 12-nm Au tags leads to a greater than 10-fold increase in angle shift, corresponding to a more than 1000-fold improvement in sensitivity for the target oligonucleotide as compared to the unamplified binding event. The extremely large angle shifts obsd. in particle-amplified SPR make it possible to conduct SPR imaging expts. on \*\*\*DNA\*\*\*

macroarrays with picomolar detection limits. A second application of metal \*\*\*nanoparticles\*\*\* employs striped multimetal rod-shaped particles as identifiable supports for \*\*\*hybridization\*\*\* assays. The striping pattern can be read via reflectivity in an optical microscope, and is used to identify the chem. on the surface of the rod. Because of the large no. of striping patterns that can be prep'd., these nanoscale "barcodes" are attractive for multiplexed bioassays.

L6 ANSWER 47 OF 63 CAPLUS COPYRIGHT 2006 ACS on STN AN 2001:197149 CAPLUS

TI Enhanced atomic force microscopy imaging of \*\*\*DNA\*\*\* \*\*\*hybridization\*\*\* by utilizing \*\*\*DNA\*\*\* -capped gold \*\*\*nanoparticles\*\*\*

AU Han, Shubo; Lin, Jianqiao; Zhou, Feimeng

CS Chemistry and Biochemistry, California State University, Los Angeles, Los Angeles, CA, 90032, USA

SO Abstracts of Papers, 221st ACS National Meeting, San Diego, CA, United States, April 1-5, 2001 (2001) ANYL-063 CODEN: 69FZD4

PB American Chemical Society

DT Journal; Meeting Abstract

LA English

AB A novel assay for selective detn. of polynucleotides using AFM together with the formation of the probe-target-\*\*\*DNA\*\*\* /gold nanoparticle sandwich structure at gold surface is described. A 17mer probe was attached to the surface for subsequent hybridization with a polynucleotide target. The hybridization efficiency can only be estd. to be about 1.1% since certain surface features could not be resolved. The 30mer-capped gold nanoparticles, not only provides another dimension of selectivity, but also reoriented the previously formed probe-target hybrid in such a way that the strands of the target become tethered with respect to the surface. Due to the improvement in resolving the hybridized target mols., an accurate detn. of the hybridization efficiency (16.5%) was achieved. Quartz crystal microbalance (QCM) was also employed to quantify the hybridization and the results are compared with the findings of AFM.

L6 ANSWER 48 OF 63 CAPLUS COPYRIGHT 2006 ACS on STN AN 2001:100554 CAPLUS

DN 134:262964

TI Site-selective immobilization of gold nanoparticles functionalized with \*\*\*DNA\*\*\* oligomers

AU Niemeyer, C. M.; Ceyhan, B.; Gao, S.; Chi, L.; Peschel, S.; Simon, U.

CS FB2 - UFT Biotechnologie und Molekulare Genetik, Universitat Bremen, Bremen, 28359, Germany

SO Colloid and Polymer Science (2001), 279(1), 68-72 CODEN: CPMSB6; ISSN: 0303-402X

PB Springer-Verlag

DT Journal

LA English

AB The organization of metal and semiconductor nanoparticles to form micro- and nanostructured assemblies is currently of tremendous interest. This communication reports on the utilization of \*\*\*DNA\*\*\* mols. as positioning elements for generating microstructured surface architecture from gold nanoparticles. Citrate-passivated 40 nm gold colloids were modified by chemisorptive coupling with a 5'-thiol-derivatized \*\*\*DNA\*\*\* oligomer. The nucleic acid was used as a mol. handle for the specific immobilization on solid supports, previously functionalized with capture \*\*\*DNA\*\*\* oligomers, complementary to the nanoparticle-bound \*\*\*DNA\*\*\*. As a consequence of the enormous specificity of nucleic acid

\*\*\*hybridization\*\*\*, the \*\*\*DNA\*\*\* -directed immobilization (DDI) allows, to site-specifically target the hybrid \*\*\*nanoparticles\*\*\* to microlocations which contain the complementary oligomers. The site-selectivity of the surface adsorption is demonstrated by immobilizing the gold colloids on a \*\*\*DNA\*\*\* microarray on a glass cover slide. Moreover, scanning force microscopy (SFM) anal., used to characterize the intermediate steps of the DDI on a gold substrate, provided initial insights into the specificity and efficiency of this technique. The application of the DDI to fabricate complex colloidal micro- and nanostructures is anticipated.

RE.CNT 21 THERE ARE 21 CITED REFERENCES AVAILABLE FOR THIS RECORD ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 49 OF 63 CAPLUS COPYRIGHT 2006 ACS on STN AN 2000:855382 CAPLUS

DN 134:173566

TI Oligonucleotide-Capped Gold Nanoparticles for Improved Atomic Force Microscopic Imaging and Enhanced Selectivity in Polynucleotide Detection

AU Han, Shubo; Lin, Jianqiao; Zhou, Feimeng; Vellanoweth, Robert Luis

CS Department of Chemistry and Biochemistry, California State University at Los Angeles, Los Angeles, CA, 90032, USA

SO Biochemical and Biophysical Research Communications (2000), 279, 265-269 CODEN: BBRC99; ISSN: 0006-291X

PB Academic Press

DT Journal

LA English

AB A novel assay for selective detn. of polynucleotides using at. force microscopy in conjunction with the formation of the probe/target/ \*\*\*DNA\*\*\* -gold nanoparticle sandwich structure at a gold surface is described. A 17-mer probe was attached to the surface for subsequent hybridization with a polynucleotide target. Due to the flat orientation of the probe-target hybrid with respect to the surface and the spatial obstruction of the unhybridized probes near the hybrids, the AFM images are not clear. The hybridization efficiency was estd. to be about 1.1% since certain surface features could not be resolved. The utilization of 30-mer-capped gold nanoparticles not only provides another dimension of selectivity, but also reorients the previously formed probe-target hybrid in such a way that the strands of the target become tethered with respect to the surface. This reorientation improves the resoln. in imaging the hybridized target mols. and provides an accurate detn. of the hybridization efficiency (16%). (c) 2000 Academic Press.

RE.CNT 37 THERE ARE 37 CITED REFERENCES AVAILABLE FOR THIS RECORD ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 50 OF 63 CAPLUS COPYRIGHT 2006 ACS on STN AN 2000:794503 CAPLUS

TI Kinetic control of oligonucleotide hybridization in monolayer nucleic acid films measured by in situ quantitative surface plasmon resonance spectroscopy.

AU Georgiadis, Rosina M.; Peterson, Alexander W.; Heaton, Richard H.

CS Department of Chemistry, Boston University, Boston, MA, 02215, USA

SO Abstracts of Papers, 220th ACS National Meeting, Washington, DC, United States, August 20-24, 2000 (2000)

COLL-311 CODEN: 69FZC3

PB American Chemical Society

DT Journal; Meeting Abstract

LA English



AB Few exptl. studies have measured the rates of oligonucleotide \*\*\*hybridization\*\*\* at surfaces despite the importance of these measurements in a range of areas including research on \*\*\*DNA\*\*\* -directed assembly of \*\*\*nanoparticles\*\*\*, \*\*\*DNA\*\*\* chip development and other biosensor applications. We present quant. SPR spectroscopy measurements of the kinetics of hybridization at surface immobilized monolayer films for a variety of oligonucleotide duplexes of varying length and base-pair sequence. We find that the rate of hybridization depends strongly not only on the degree of mismatch in the base-pair sequence but also on the position at which the duplex forms along the immobilized strand. The kinetics of surface duplex formation are much slower than rates which have been measured in homogenous aq. soln. Melting curves for surface immobilized duplex dehybridization, measured by temp. dependent SPR spectroscopy are compared with results in homogenous soln. as measured by CD spectroscopy. Our results are consistent with kinetic control of the rate of duplex formation in immobilized monolayer nucleic acid films.

L6 ANSWER 51 OF 63 CAPLUS COPYRIGHT 2006 ACS on STN  
AN 2000:792856 CAPLUS  
DN 134:97279  
TI Chip-Based Optical Detection of \*\*\*DNA\*\*\* Hybridization by Means of Nanobead Labeling  
AU Reichert, Joerg; Csaki, Andrea; Koehler, J. Michael; Fritzsche, Wolfgang  
CS Institute of Physical High Technology, Jena, D-07702, Germany  
SO Analytical Chemistry (2000), 72(24), 6025-6029 CODEN: ANCHAM; ISSN: 0003-2700  
PB American Chemical Society  
DT Journal  
LA English  
AB A new scheme for the detection of mol. interactions based on optical readout of nanoparticle labels has been developed. Capture \*\*\*DNA\*\*\* probes were arrayed on a glass chip and incubated with nanoparticle-labeled target \*\*\*DNA\*\*\* probes, contg. a complementary sequence. Binding events were monitored by optical means, using reflected and transmitted light for the detection of surface-bound nanoparticles. Control expts. exclude significant influence of nonspecific binding on the obsd. contrast. Scanning force microscopy revealed the distribution of nanoparticles on the chip surface.  
RE.CNT 12 THERE ARE 12 CITED REFERENCES AVAILABLE FOR THIS RECORD ALL CITATIONS AVAILABLE IN THE REFORMAT

L6 ANSWER 52 OF 63 CAPLUS COPYRIGHT 2006 ACS on STN  
AN 2000:683718 CAPLUS  
DN 135:895  
TI Facile colorimetric detection of polynucleotides based on gold nanoparticle probes  
AU Storhoff, James J.; Elghanian, Robert; Mucic, Robert C.; Mirkin, Chad A.; Letsinger, Robert L.  
CS Department of Chemistry, Northwestern University, Evanston, IL, 60208, USA  
SO Proceedings of the ERDEC Scientific Conference on Chemical and Biological Defense Research, Aberdeen Proving Ground, MD, United States, Nov. 17-20, 1998 (1999), Meeting Date 1998, 221-226. Editor(s): Berg, Dorothy A. Publisher: National Technical Information Service, Springfield, Va. CODEN: 69AJH3  
DT Conference  
LA English

AB Novel probes for polynucleotide detection can be prepd. by functionalizing gold nanoparticles with alkythiol-modified oligonucleotides. These probes readily \*\*\*hybridize\*\*\* to complementary polynucleotides in soln. leading to the formation of a polymeric network of gold \*\*\*nanoparticles\*\*\*. The particles serve as reporter groups, triggering a red to pinkish/purple color change upon hybridization. By spotting the test solns. onto a solid-phase support, target polynucleotides can be detected colorimetrically and differentiated from targets contg. single base imperfections.  
RE.CNT 18 THERE ARE 18 CITED REFERENCES AVAILABLE FOR THIS RECORD ALL CITATIONS AVAILABLE IN THE REFORMAT

L6 ANSWER 53 OF 63 CAPLUS COPYRIGHT 2006 ACS on STN  
AN 2000:624265 CAPLUS  
DN 134:111002  
TI Colloidal Au-enhanced surface plasmon resonance for ultrasensitive detection of \*\*\*DNA\*\*\* hybridization  
AU He, Lin; Musick, Michael D.; Nicewarner, Sheila R.; Salinas, Frank G.; Benkovic, Stephen J.; Natan, Michael J.; Keating, Christine D.  
CS Department of Chemistry, The Pennsylvania State University, University Park, PA, 16802, USA  
SO Journal of the American Chemical Society (2000), 122(38), 9071-9077 CODEN: JACSAT; ISSN: 0002-7863  
PB American Chemical Society  
DT Journal  
LA English  
AB A new approach to ultrasensitive detection of \*\*\*DNA\*\*\* \*\*\*hybridization\*\*\* based on \*\*\*nanoparticle\*\*\* -amplified surface plasmon resonance (SPR) is described. Use of the Au nanoparticle tags leads to a greater than 10-fold increase in angle shift, corresponding to a more than 1000-fold improvement in sensitivity for the target oligonucleotide as compared to the unamplified binding event. This enhanced shift in SPR reflectivity is a combined result of greatly increased surface mass, high dielec. const. of Au particles, and electromagnetic coupling between Au nanoparticles and the Au film. \*\*\*DNA\*\*\* melting and digestion expts. further supported the feasibility of this approach in \*\*\*DNA\*\*\* hybridization studies. The extremely large angle shifts obsd. in particle-amplified SPR make it possible to conduct SPR imaging expts. on \*\*\*DNA\*\*\* arrays. In the present work, macroscopic 4 .times. 4 arrays were employed, and a .apprx.10 pM limit of quantitation was achieved for 24-mer oligonucleotides (surface d. .ltoreq.8 .times. 108 mols./cm<sup>2</sup>). Even without further optimization, the sensitivity of this technique begins to approach that of traditional fluorescence-based methods for \*\*\*DNA\*\*\* hybridization. These results illustrate the potential of particle-amplified SPR for array-based \*\*\*DNA\*\*\* anal. and ultrasensitive detection of oligonucleotides.  
RE.CNT 64 THERE ARE 64 CITED REFERENCES AVAILABLE FOR THIS RECORD ALL CITATIONS AVAILABLE IN THE REFORMAT

L6 ANSWER 54 OF 63 CAPLUS COPYRIGHT 2006 ACS on STN  
AN 2000:403224 CAPLUS  
DN 133:219622  
TI The \*\*\*DNA\*\*\* -Mediated Formation of Supramolecular Mono- and Multilayered Nanoparticle Structures  
AU Taton, T. Andrew; Mucic, Robert C.; Mirkin, Chad A.; Letsinger, Robert L.  
CS Department of Chemistry and Center for Nanofabrication and Molecular Self-Assembly, Northwestern University, Evanston, IL, 60208, USA



SO Journal of the American Chemical Society (2000), 122(26), 6305-6306 CODEN: JACSAT; ISSN: 0002-7863

PB American Chemical Society

DT Journal

LA English

AB Herein, we report a new, noncovalent \*\*\*DNA\*\*\* -based strategy for controlling the stepwise growth of layered nanoparticle structures off of glass surfaces. This is the first reported methodol. for forming layered \*\*\*nanoparticle\*\*\* -based materials from \*\*\*DNA\*\*\* and designed \*\*\*hybridization\*\*\* events. The strategy should be quite general and extendable to other nanoparticle sizes and compns. The unexpected sharp melting properties of single particles hybridized to the glass support point to a possible route to developing chip-based detection formats for \*\*\*DNA\*\*\* with selectivities that are better than those of conventional probe systems.

RE.CNT 18 THERE ARE 18 CITED REFERENCES AVAILABLE FOR THIS RECORD ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 55 OF 63 CAPLUS COPYRIGHT 2006 ACS on STN AN 2000:350693 CAPLUS DN 134:37640

TI Amplified microgravimetric gene sensor using Au

nanoparticle modified oligonucleotides

AU Zhou, Xi Chun; O'Shea, Sean J.; Li, Sam Fong Yau

CS Institute of Materials Research and Engineering, 117602, Singapore

SO Chemical Communications (Cambridge) (2000), (11), 953-954 CODEN: CHCOFS; ISSN: 1359-7345

PB Royal Society of Chemistry

DT Journal

LA English

AB A novel microgravimetric gene sensing system has been developed using an oligonucleotide anchored on the gold electrode of a quartz crystal microbalance and an Au nanoparticle modified oligonucleotide, both of which formed a sandwich-type ternary complex with the target \*\*\*DNA\*\*\* to give an amplified frequency response.

RE.CNT 11 THERE ARE 11 CITED REFERENCES AVAILABLE FOR THIS RECORD ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 56 OF 63 CAPLUS COPYRIGHT 2006 ACS on STN AN 2000:345850 CAPLUS DN 133:147119

TI A gold nanoparticle/latex microsphere-based colorimetric oligonucleotide detection method

AU Reynolds, Robert A., III; Mirkin, Chad A.; Letsinger, Robert L.

CS Department of Chemistry and Center for Nanofabrication and Molecular Self-Assembly, Northwestern University, Evanston, IL, 60208, USA

SO Pure and Applied Chemistry (2000), 72(1-2), 229-235 CODEN: PACHAS; ISSN: 0033-4545

PB International Union of Pure and Applied Chemistry

DT Journal

LA English

AB An exceptionally simple and effective \*\*\*DNA\*\*\* detection methodol. based on latex microsphere and gold nanoparticle probes has been developed. The latex and gold particle probes, which were functionalized with sep. oligonucleotide sequences, undergo hybridization in the presence of target strands that are complementary to both of the probes. Duplex formation thus results in linking of gold nanoparticles to

the latex microspheres and a corresponding white-to-red color change, which, because of the particularly large extinction coeff. of the gold nanoparticles, is clearly visible to the naked eye.

Background signal caused by unbound gold nanoparticles is significantly reduced by filtering the soln. contg. the sample and probes through a size-selective cellulose acetate membrane. The unbound gold probes move freely through this membrane while the larger latex particles are trapped. Therefore, if the latex and gold nanoparticles are joined together via the target oligonucleotides, the membrane appears red, indicating a pos. test result. If no hybridization takes place, the membrane appears white, indicating a neg. result. The lower detection limits for this system are 500 pM for a 24 base single-stranded target and 2.5 nM for a duplex target oligonucleotide.

RE.CNT 12 THERE ARE 12 CITED REFERENCES AVAILABLE FOR THIS RECORD ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 57 OF 63 CAPLUS COPYRIGHT 2006 ACS on STN AN 2000:335606 CAPLUS DN 133:2215

TI Signal amplification by crosslinking of \*\*\*nanoparticles\*\*\* by \*\*\*hybridization\*\*\* of bound nucleic acids

IN Barbera-Guillem, Emilio; Nelson, M. Bud; Castro, Stephanie PA Biocrystal Limited, USA

SO PCT Int. Appl., 72 pp. CODEN: PIXXD2

DT Patent

LA English

FAN.CNT	12	PATENT NO.	KIND	DATE	APPLICATION NO.
NO.	DATE	-----	---	-----	-----

PI WO 2000028088 A1 20000518 WO 1999-US26612 19991110 W: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM RW: GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG US 6261779 B1 20010717 US 1999-437076

19991109 WO 2000028089 A1 20000518 WO 1999-US26616 19991110 W: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM RW: GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG EP 1157133 A1 20011128 EP 1999-960265

19991110 R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO PRAI US 1998-107828P P 19981110 US 1999-437076 A 19991109 US 1998-109626P P 19981124 WO 1999-US26612 W 19991110

AB A method of amplifying a signal by crosslinking of water-sol. nanocrystals using nucleic acids immobilized on the particles is described. An anal. probe, e.g. an antibody, attached to one type of particle carrying several mols. of one nucleic acid sequence is used to bind the analyte. This is then incubated with a second type of particle carrying a probe complementary to the nucleic acid on the first type of particle. By alternating incubations with the two types of particles, large branching

complexes are built up and easily detected, e.g by fluorometry using a nucleic acid binding dye, or by use of fluorescent nanocrystals (quantum dots). Methods of prepg. quantum dots to stabilize them against oxidn. by capping unused reactive groups on the surface are described.

RE.CNT 3 THERE ARE 3 CITED REFERENCES AVAILABLE FOR THIS RECORD ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 58 OF 63 CAPLUS COPYRIGHT 2006 ACS on STN  
AN 2000:333894 CAPLUS  
DN 133:130412

TI Molecular beacon biosensors for \*\*\*DNA\*\*\* / \*\*\*RNA\*\*\* analysis

AU Fang, Xiaohong; Schuster, Sheldon; Liu, Xiaojing; Correll, Tiffany; Zhang, Peng; Tapecc, Ruby; Santra, Swadeshmukul; Qhobosheanne, Monde; Lou, Jane Hua; Tan, Weihong  
CS UF Brain Institute, Dep. Chem., Univ. Florida, Gainesville, FL, USA

SO Proceedings of SPIE-The International Society for Optical Engineering (2000), 3926(Advances in Nucleic Acid and Protein Analyses, Manipulation, and Sequencing), 2-8 CODEN: PSISDG; ISSN: 0277-786X

PB SPIE-The International Society for Optical Engineering  
DT Journal  
LA English

AB We have developed a variety of novel \*\*\*DNA\*\*\* biosensors using a new class of oligonucleotide probe, mol. beacon (MB). MB has the fluorescence signal transduction mechanism built within the mols. It can report the presence of specific nucleic acids with high sensitivity and excellent selectivity. Biotinylated MBs have been designed and synthesized for immobilization onto silica surface through avidin-biotin binding. The effect of the avidin-biotin bridge on the MB hybridization has been studied. Our result shows that using streptavidin has less effect than using avidin in MB hybridization. Two kinds of fiber optical \*\*\*DNA\*\*\* sensors have been prepd. and characterized: a fiber optic evanescent wave sensor and a submicrometer optical fiber sensor. The sensors are rapid, stable, highly selective, reproducible and regenerable. They have been applied to detect specific \*\*\*DNA\*\*\* and \*\*\*mRNA\*\*\* sequences and to the study of the \*\*\*DNA\*\*\* hybridization kinetics. Silica nanoparticles have also been used for MB immobilization in order to prep. a large quantity of nanometer sized \*\*\*DNA\*\*\* / \*\*\*RNA\*\*\* biosensors.

RE.CNT 21 THERE ARE 21 CITED REFERENCES AVAILABLE FOR THIS RECORD ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 59 OF 63 CAPLUS COPYRIGHT 2006 ACS on STN  
AN 2000:331439 CAPLUS

TI Fluorescence-based method for the determination of surface coverage and \*\*\*hybridization\*\*\* efficiency of thiol-capped oligonucleotides bound to gold \*\*\*nanoparticles\*\*\*.

AU Demers, Linette M.; Mucic, Robert C.; Reynolds, Robert A., III; Mirkin, Chad A.; Letsinger, Robert L.  
CS Department of Chemistry and Center for Nanofabrication and Molecular Self-assembly, Northwestern University, Evanston, IL, 60208, USA

SO Book of Abstracts, 219th ACS National Meeting, San Francisco, CA, March 26-30, 2000 (2000), INOR-649 Publisher: American Chemical Society, Washington, D. C. CODEN: 69CLAC  
DT Conference; Meeting Abstract  
LA English

AB Recently, our group reported a new \*\*\*DNA\*\*\* detection scheme based upon the \*\*\*hybridization\*\*\* of target

\*\*\*DNA\*\*\* to gold \*\*\*nanoparticle\*\*\* probes with surface immobilized alkylthiol-capped oligonucleotides. The basis for this detection method is a colorimetric change assocd. with particle aggregation caused by oligonucleotide hybridization events at the surface. Two factors that dictate the magnitude of the observable optical changes are: 1) the extent of surface modification with the oligonucleotides, and 2) the ability of complementary oligonucleotides to access surface-bound oligonucleotides. In this work we describe a fluorescence-based method for detg. the surface coverage of the \*\*\*nanoparticle\*\*\* -bound oligonucleotides, and the efficiency of \*\*\*hybridization\*\*\* with complementary oligonucleotides. Importantly, this method has enabled us to investigate parameters such as electrolyte concn., oligonucleotide sequence and length, and the role of diluent mols., which influence the surface coverage and resulting \*\*\*hybridization\*\*\* properties of oligonucleotide-functionalized \*\*\*nanoparticles\*\*\*. In addn., we show that variation of the loading conditions affords control over the \*\*\*hybridization\*\*\* efficiency of the oligonucleotide-modified \*\*\*nanoparticles\*\*\*.

L6 ANSWER 60 OF 63 CAPLUS COPYRIGHT 2006 ACS on STN  
AN 1999:703246 CAPLUS  
DN 132:31330

TI Chemistry of oligonucleotide-gold nanoparticle conjugates

AU Letsinger, Robert L.; Mirkin, Chad A.; Elghanian, Robert; Mucic, Robert C.; Storhoff, James J.  
CS Department of Chemistry, Northwestern University, Evanston, IL, 60208, USA

SO Phosphorus, Sulfur and Silicon and the Related Elements (1999), 144-146, 359-362 CODEN: PSSLEC; ISSN: 1042-6507  
PB Gordon & Breach Science Publishers

DT Journal; General Review  
LA English

AB A review, with 10 refs. Conjugates prepd. by immobilizing thiol-terminated oligonucleotides onto gold nanoparticles form stable colloidal solns. in aq. media. The oligonucleotides can serve as linkers to organize the gold particles reversibly into three dimensional assemblies, and the gold particles can function as colorimetric reporters for hybridization of the bound oligomers with target oligonucleotides in soln.

RE.CNT 13 THERE ARE 13 CITED REFERENCES AVAILABLE FOR THIS RECORD ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 61 OF 63 CAPLUS COPYRIGHT 2006 ACS on STN  
AN 1998:150275 CAPLUS  
DN 128:226756

TI One-Pot Colorimetric Differentiation of Polynucleotides with Single Base Imperfections Using Gold Nanoparticle Probes

AU Storhoff, James J.; Elghanian, Robert; Mucic, Robert C.; Mirkin, Chad A.; Letsinger, Robert L.  
CS Department of Chemistry, Northwestern University, Evanston, IL, 60208, USA

SO Journal of the American Chemical Society (1998), 120(9), 1959-1964 CODEN: JACSAT; ISSN: 0002-7863

PB American Chemical Society  
DT Journal  
LA English

AB Selective colorimetric polynucleotide detection based on Au nanoparticle probes which align in a "tail-to-tail" fashion onto a target polynucleotide is described. In this new nanoparticle-based detection system, Au particles (.apprx.13 nm diam.), which are capped with 3'- and 5'-(alkanethiol)oligonucleotides, are used to complex a 24-base polynucleotide target.

\*\*\*Hybridization\*\*\* of the target with the probes results in the

formation of an extended polymeric Au \*\*\*nanoparticle\*\*\* /polynucleotide aggregate, which triggers a red to purple color change in soln. The color change is due to a red shift in the surface plasmon resonance of the Au nanoparticles. The aggregates exhibit characteristic, exceptionally sharp "melting transitions" (monitored at 260 or 700 nm), which allows one to distinguish target sequences that contain one base end mismatches, deletions, or an insertion from the fully complementary target. When test solns. are spotted onto a C18 reverse-phase thin-layer chromatog. plate, color differentiation is enhanced and a permanent record of the test is obtained, thereby providing a better method for distinguishing the aforementioned target sequences. Significantly, one-pot colorimetric detection of the target in the presence of four strands with single base imperfections can be accomplished with this new probe system.

RE.CNT 23 THERE ARE 23 CITED REFERENCES AVAILABLE FOR THIS RECORD ALL CITATIONS AVAILABLE IN THE RE FORMAT

L6 ANSWER 62 OF 63 CAPLUS COPYRIGHT 2006 ACS on STN AN 1997:458356 CAPLUS DN 127:140241  
TI Modification and immobilization of enzymes and other biological macromolecules by polymer materials and their biomedical applications  
AU Ma, Jian-Biao  
CS State Key Laboratory Functional Polymer Materials Adsorption Separation, Institute Polymer Chemistry, Nankai University, Tianjin, 300071, Peop. Rep. China  
SO Gaodeng Xuexiao Huaxue Xuebao (1997), 18(7), 1226-1235 CODEN: KTHPDM; ISSN: 0251-0790  
PB Gaodeng Jiaoyu Chubanshe  
DT Journal; General Review  
LA Chinese  
AB A review with 30 refs. Recent progress in biomedical polymers, chem. modification of L-asparaginase by natural and synthetic polymers, synthesis of dextrin magnetic \*\*\*nanoparticles\*\*\* as carriers during the immobilization of enzyme and antibodies, polymeric latexes useful in the detection of \*\*\*DNA\*\*\* \*\*\*hybridization\*\*\*, enzyme immobilization on porous polymeric beads, immobilization of enzymes in the conducting polymer membrane and fabrication of bioelectrodes, as well as biomimetic polymers of mol. recognition.

L6 ANSWER 63 OF 63 CAPLUS COPYRIGHT 2006 ACS on STN AN 1996:349849 CAPLUS DN 125:41765  
TI Adsorption of antisense oligonucleotides onto nanoparticles for therapeutic use.  
IN Helene, Claude; Saison, Behmoaras Ester  
PA Centre National De La Recherche Scientifique Cnrs, Fr.  
SO Fr. Demande, 23 pp. CODEN: FRXXBL  
DT Patent  
LA French

FAN.CNT 1 PATENT NO.	KIND	DATE	APPLICATION
NO.	DATE		
PI FR 2724935	A1	19960329	FR 1994-11512
19940927 FR 2724935	B1	19961220	
PRAI FR 1994-11512		19940927	

AB Methods for adsorption or encapsulation of antisense oligonucleotides onto nanoparticles for use in the treatment of tumors or other diseases assocd. with disruption of patterns of regulation of gene expression are described. Adsorption onto \*\*\*nanoparticles\*\*\* increases the resistance of the

oligonucleotides to nucleases and improved cellular uptake and \*\*\*hybridization\*\*\* to the target sequence. Specifically, antisense oligonucleotides directed against mutants of the ras oncogene are described. Adsorbed or immobilized oligonucleotides are up to 100-fold more effective than the free oligonucleotide. The nanoparticles have a diam. of 50-500 nm and include a hydrophobic cationic polymer for efficient binding of the oligonucleotides. Nude mice inoculated with HBL100ras cells showed strong inhibition of tumor growth when treated with nanoparticles contg. the appropriate antisense \*\*\*DNA\*\*\*.

=> log y	SINCE FILE	
COST IN U.S. DOLLARS	ENTRY	SESSION
TOTAL	202.46	202.67
FULL ESTIMATED COST		

DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)	SINCE	
FILE TOTAL	ENTRY	
SESSION		
CA SUBSCRIBER PRICE	-45.75	-45.75

STN INTERNATIONAL LOGOFF AT 17:53:07 ON 27 JAN 2006